

Konrad-Zuse-Zentrum für Informationstechnik Berlin

Homoclinic Bifurcation at Resonant Eigenvalues

Shui-Nee Chow[†] Bo Deng^{††} Bernold Fiedler^{†††,*}

- School of Mathematics Georgia Institute of Technology Atlanta, Georgia 30332 U.S.A.
- Dept. of Mathematics and Statistics University of Nebraska Lincoln, NE 68588 - 0323 U.S.A.
- Institute of Applied Mathematics University of Heidelberg
 Im Neuenheimer Feld 294
 D- 6900 Heidelberg
 Federal Republic of Germany
 - * Partially supported by the Deutsche Forschungsgemeinschaft and by Konrad-Zuse-Zentrum für Informationstechnik Berlin

Preprint SC 88-10 (December 1988)

Contents

1.	Introduction	1	
2.	A model example: the piecewise linear case	12	
3.	Shilnikov variables and the strong λ -Lemma	17	
4.	Ljapunov - Schmidt reduction	23	
5.	The side-switching bifurcation	34	
6.	The homoclinic doubling bifurcation	39	
7.	Discussion	52	
8.	Appendix: genericity assumptions	65	
A	Acknowledgement		
Re	References		

1. Introduction

By definition, a homoclinic orbit $z^*(t)$ of a flow

$$\dot{z} = F(z) \quad , \quad z \in \mathbb{R}^{m+n} \tag{1.1}$$

tends to the same equilibrium, say z = 0, F(0) = 0, for both $t \to \pm \infty$. Homoclinic orbits are a key phenomenon for understanding more complicated dynamics, including chaotic motions; see e.g. [GUCKENHEIMER & HOLMES 1, SPARROW]. For typical vector fields F, however, homoclinic orbits do not exist. The reason is the following. Suppose z = 0 is a hyperbolic equilibrium, i.e. the eigenvalues of F'(0) stay away from the imaginary axis. Then the *m*-dimensional stable manifold W^s of z = 0 consists of all those $z_0 = z(0)$ for which

$$\lim_{t\to+\infty} z^*(t) = 0 \;\; ,$$

see e.g. [CHOW & HALE]. Likewise, the *n*-dimensional unstable manifold W^{u} is associated to

$$\lim_{t\to-\infty}z^*(t)=0.$$

By definition, W^u and W^s intersect nontrivially (i.e. at some point $\mathbf{p} = z^*(0) \neq 0$) if $z^*(t)$ is a homoclinic orbit. Note that W^u and W^s then intersect all along the orbit $z^*(t)$. In particular, the tangent spaces $T_{\mathbf{p}}W^s$ and $T_{\mathbf{p}}W^u$ intersect nontrivially,

$$\dot{z}(t) \in T_{\mathbf{p}} W^{s} \cap T_{\mathbf{p}} W^{u} \quad \text{at } t = 0.$$

$$(1.2)$$

In other words: W^{u} and W^{s} intersect nontransversely at p; and

$$\operatorname{codim}\left(T_{\mathbf{p}}W^{s} + T_{\mathbf{p}}W^{u}\right) = 1 \tag{1.3}$$

or, even worse, > 1. Perturbing the vector field F slightly, we can therefore push W^u slightly in the direction of that remaining codimension, keeping W^s fixed. Then W^u and W^s do not intersect anymore, locally, and the homoclinic orbit has disappeared. A precise statement of this idea is the Kupka-Smale theorem [KUPKA, SMALE]. This theorem asserts that for generic $F \in C^{\infty}$, stable and unstable manifolds intersect transversely, if they intersect at all. By (1.3), this excludes the possibility of homoclinic orbits. The word "generic" indicates that the theorem may fail (and it does!) only for a subset of $F \in C^{\infty}$, which is of the first Baire category in the weak Whitney topology. In particular, the assertion of the Kupka-Smale theorem holds for a dense set of F. See e.g. [ABRAHAM & ROBBIN] for more details. Note that this situation contrasts markedly with the case of discrete time flows viz. of iterating diffeomorphisms, where transverse homoclinic points can occur — leading to Smale horseshoes and shift dynamics; see e.g. [MOSER].

We have seen above, heuristically, how homoclinic orbits are a codimension 1 phenomenon in the space of vector fields. Therefore, we now consider C^{∞} parametrized vector fields.

$$\dot{z} = F(\alpha, z) \quad , \quad z \in \mathbb{R}^{m+n}$$
 (1.1_{\alpha})

with parameter α . Below, we will consider parameters $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$. But for the moment, let us discuss the case $\alpha \in \mathbb{R}$. By the above reasoning, homoclinic orbits will typically, i.e. generically, occur for isolated parameter values, say at $\alpha = 0$. What happens for small $\alpha \neq 0$? One possible scenario is the following. For $\alpha < 0$ we see a periodic orbit $z_{\alpha}(t)$ of minimal period T_{α} . For $\alpha \nearrow 0$, some part of the periodic orbits approaches the equilibrium z = 0, e.g.

$$\lim_{\alpha \nearrow 0} z_{\alpha}(0) = 0$$

while the remaining points approach the closure of the homoclinic orbit of

$$\dot{z}=F(0,\,z)$$

Simultaneously, the minimal periods T_{α} tend to infinity. For $\alpha > 0$, both the periodic orbits and the homoclinic orbit just disappear. This was termed a "blue sky catastrophe" in [ABRAHAM & MARSDEN]. For $z \in \mathbb{R}^{m+n} = \mathbb{R}^2$, and under certain additional assumptions also for m+n > 2, SHILNIKOV has shown that this description holds true for generic one parameter families F. This work dates back to 1962; see e.g. the survey [SHILNIKOV 5] (1968).

From now on we consider generic two parameter vector fields

$$F(\alpha, \cdot), \ \alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$$

Since homoclinic orbits are codimension one objects, we expect them to occur along one-dimensional curves $\alpha = \alpha(\tau)$ in two parameter space $\alpha \in \mathbb{R}^2$. This point of view makes homoclinic orbits amenable to a pathfollowing approach. We may now follow curves of homoclinic orbits rather than hitting them "catastrophically", out of the blue.

The following illustrative example goes back to [ARNOLD, BOGDANOV 1-2] (1972, 1976) and to [TAKENS] (1974):

$$\dot{z_1} = z_2$$

$$\dot{z_2} = \alpha_1 + \alpha_2 z_1 + z_1^2 + \sigma z_1 z_2 ,$$
(1.4)

where $\sigma = \pm 1$. In figure 1.1 we show the local bifurcation diagram of system (1.4).



Figure 1.1: A B-point

To the left of the fold line, two equilibria co-exist. Crossing the fold to the right, they merge and disappear. Crossing the Hopf line, one of the two equilibria undergoes a Hopf bifurcation, giving rise to periodic orbits when we decrease α_1 . The periodic orbits terminate by a blue sky catastrophe at homoclinic orbits, which occur along a one-dimensional curve of parameters α . Following that homoclinic curve towards $\alpha = 0$, the homoclinic orbits shrink down to the equilibrium z = 0. More generally, figure 1.4 describes the generic local bifurcation diagram associated to an equilibrium (α_0 , z_0) of (1.1_{α}) where $D_z F(\alpha_0, z_0)$ has an algebraically double eigenvalue 0. These results are due to [ARNOLD, BOGDANOV 1-2], (1972, 1976).

The term *B*-point was used in [FIEDLER 2] for equilibria like (α_0, z_0) . An index for *B*-points seemed to indicate a possibility for some global pathfollowing results on homoclinic orbits, cf. [FIEDLER 2, p.74]. Indeed, *B*-points give rise to branches of homoclinic orbits in generic two parameter flows just as Hopf bifurcations give rise to branches of periodic orbits in generic one parameter flows. It is then a necessary first step, towards global results on homoclinic orbits, to understand local bifurcations which a branch of homoclinic orbits can undergo on its way. We will resume this aspect at the end of the discussion in section 7. We now specify two such bifurcations: the resonant side-switching and the resonant homoclinic doubling (see Theorem A and Theorem B below). We fix some notations and some assumptions. First of all, we assume

$$F : |\mathbb{R}^2 \times |\mathbb{R}^{m+n} \to |\mathbb{R}^{m+n}$$

$$(\alpha, z) \to F(\alpha, z)$$
is a generic C^{M+5} vector field with
two parameters $\alpha = (\alpha_1, \alpha_2) \in |\mathbb{R}^2, 2 \le M \le \infty$.
$$(1.5)$$

As before, genericity means that our results will hold for a residual subset of $C^{M+5}(\mathbb{R}^2 \times \mathbb{R}^{m+n}, \mathbb{R}^{m+n})$ in the weak Whitney topology of C^k -convergence, i.e., uniformly for compact sets and bounded sets of $k \leq M+5$.

Our second assumption concerns the existence of a homoclinic orbit Γ with real principal eigenvalues. We assume

the vector field $F(0, \cdot)$ at $\alpha = 0$ admits a homoclinic orbit

 $\Gamma = \{z^*(t) : t \in \mathbb{R}\}$

such that

$$\lim_{t\to\pm\infty}z^*(t)=0\;.$$

Furthermore, the linearization $D_z F(0, 0)$ at the equilibrium (1.6)

z = 0 has simple real eigenvalues $-\mu_0 < 0 < \nu_0$ such that

any remaining eigenvalue μ of $D_z F(0, 0)$ satisfies

either Re $\mu < -\bar{\mu_1} < -\mu_0 < 0$ or Re $\nu > \bar{\nu_1} > \nu_0 > 0$.

We call $-\mu_0$, ν_0 the principal eigenvalues of $D_z F(0, 0)$, for the following reason. There exist submanifolds W^{ss} ("strong stable") and W^{uu} ("strong unstable") of the *m*-dimensional stable manifold W^s and the *n*-dimensional unstable manifold W^u of z = 0, respectively, such that

$$\lim_{t \to -\infty} e^{\bar{\nu}_1 |t|} z(t) = 0 \quad \text{on } W^{uu} \text{, and}$$

$$\lim_{t \to +\infty} e^{\bar{\mu}_1 t} z(t) = 0 \quad \text{on } W^{ss} \text{.}$$
(1.7)

It turns out that dim $W^{ss} = m - 1$, dim $W^{uu} = n - 1$. The tangent spaces at z = 0 are given by the parts of the spectrum of $D_z F(0, 0)$ with real part $< -\bar{\mu_1}$ and $> \bar{\nu_1}$, respectively. Moreover, for any $z(0) \in W^s \setminus W^{ss}$, the limit

$$\lim_{t \to +\infty} z(t) / |z(t)| \neq 0 \tag{1.8}$$

exists and is a unit eigenvector of the principal stable eigenvalue $-\mu_0$. An analogous statement holds for $z(0) \in W^u \setminus W^{uu}$ and the principal unstable eigenvalue ν_0 . For a reference see [HIRSCH & PUGH & SHUB, SHUB, BRUNOVSKÝ & FIEDLER 1].

We now assume the resonance condition

$$\mu_0 = \nu_0 \tag{1.9}$$

for the principal eigenvalues $-\mu_0 < 0 < \nu_0$. We remember that we expect homoclinic orbits to occur along one-dimensional curves in parameter space. Therefore, resonance condition (1.9) will not contradict our genericity assumption (1.5). Of course, here we have shifted the associated equilibrium (α_0, z_0), where (1.9) holds to (0, 0).

As a final prelude to our main results, we distinguish between twisted and nontwisted homoclinic orbits (cf. figures 1.2.a,b where m = 1, n = 2).



Figure 1.2: Homoclinic orbits

Let $\Gamma = \{z^*(t) : t \in \mathbb{R}\}$ denote a homoclinic orbit of a vector field with simple real principal eigenvalues, as in (1.6). Choose points $\mathbf{p}, \mathbf{q} \in \Gamma$ sufficiently close to the equilibrium z = 0 associated to Γ , say $\mathbf{p} = z^*(0), \mathbf{q} = z^*(T)$. We assume nondegeneracy of Γ ,

$$\operatorname{codim} (T_{\mathbf{p}}W^{u} + T_{\mathbf{p}}W^{s}) = 1$$
, (1.10.a)

in accordance with (1.3). Motivated by the convergence result (1.8), we further

assume the following general position of Γ :

$$\mathbf{p} \notin W^{ss}$$
, $\mathbf{p} \notin W^{uu}$. (1.10.b)

Obviously, assumptions (1.10.a,b) imply the same statements for q. Define the two unit vectors

$$e^{\pm} := \pm \lim_{t \to \mp \infty} \dot{z}^{*}(t) / |\dot{z}^{*}(t)| \quad . \tag{1.11}$$

By (1.10.b), $e^+ \in T_0 W^u$ is a unit eigenvector of $\nu_0 > 0$, and $e^- \in T_0 W^s$ belongs to $-\mu_0 < 0$. Now define the hyperplanes

$$T_{z^{\bullet}(t)} := T_{z^{\bullet}(t)} W^{s} + T_{z^{\bullet}(t)} W^{u} \quad . \tag{1.12}$$

Note that codim $T_{z^{\bullet}(t)} = 1$, by assumption (1.10.a). For simple real principal eigenvalues, it is now a consequence of the strong λ -Lemma, that generically

$$\lim_{t \to -\infty} T_{z^{\bullet}(t)} = T_0 W^{ss} \oplus T_0 W^u$$

$$\lim_{t \to +\infty} T_{z^{\bullet}(t)} = T_0 W^s \oplus T_0 W^{uu} .$$
(1.13)

See the strong inclination property in [DENG 1]. This is illustrated in figure 1.2, where $W^{ss} = \{0\}$ since $m = \dim W^s = 1$. For a precise statement of the strong λ -Lemma see lemma 3.3 below. Choosing $\mathbf{p}, \mathbf{q} \in \Gamma$ close enough to z = 0, as above, (1.13) implies

$$e^- \notin T_{\mathbf{p}}$$
, $\mathbb{R}^{m+n} = T_{\mathbf{p}} \oplus \operatorname{span}(e^-)$
 $e^+ \notin T_{\mathbf{q}}$, $\mathbb{R}^{m+n} = T_{\mathbf{q}} \oplus \operatorname{span}(e^+)$. (1.14)

Finally, note that $T_{z^{\bullet}(t)}$, $0 \le t \le T$, defines a homotopy from $T_{\mathbf{p}}$ to $T_{\mathbf{q}}$, justifying the following definition of a twist.

Definition: Let Γ be a nondegenerate homoclinic orbit in general position and with real principal eigenvalues, that is (1.6), (1.10.a,b), (1.13) hold.

We call Γ twisted, if e^- and e^+ point to opposite sides of T_p and T_q , respectively. See figure 1.2.b.

If e^- , e^+ point to the same side of T_p , T_q , respectively, then we call Γ non-twisted. See figure 1.2.a.

Clearly, twisted homoclinic orbits can occur only in space dimensions ≥ 3 . Homoclinic orbits in planar flows are always non-twisted. Note that the twist can also be expressed by watching the winding of the bundle of normal vectors to $T_{z^*(t)}$ as $z^*(t)$ moves from **p** to **q**. We need a final piece of terminology. Fix a small tubular neighborhood U of our (twisted or non-twisted) homoclinic orbit $clos \Gamma = \Gamma \cup \{0\}$. An *N-periodic orbit* is a periodic orbit which is contained in U and has winding number N in U. Similarly, we define an *N-homoclinic orbit*. In particular, Γ itself is a 1-homoclinic orbit. As long as U is chosen small enough, the above definition is independent of the particular choice of U. The terminology extends canonically to small perturbations of the original vector field $F(0, \cdot)$. We can now state our main results.

Theorem A: Resonant side-switching.

Let $F = F(\alpha, z)$ of class C^{M+5} , $2 \leq M \leq \infty$, be a generic two parameter vector field with a non-twisted resonant homoclinic orbit at $\alpha = 0$. Then resonant side-switching occurs at $\alpha = 0$ (see figure 1.3.a).

In more detail, let assumptions (1.5), (1.6), (1.9) be satisfied for a non-twisted homoclinic orbit Γ (cf. definition 1.1). Let U denote a sufficiently small tubular neighborhood of Γ .

Then there exist a C^{M} diffeomorphic local change of parameters

$$\epsilon = (\epsilon_1, \epsilon_2) = \epsilon(\alpha)$$

at $\alpha = 0$ and a function

$$\epsilon_{2} = \kappa(\epsilon_{1}) \begin{cases} = 0 & \epsilon_{1} \leq 0 \\ for & \\ > 0 & \epsilon_{1} > 0 \end{cases}$$
(1.15)

of class C^M for $\epsilon_1 > 0$, such that the numbers of 1-periodic (1-per) and of 1homoclinic (1-hom) orbits in U for the parameter regions 0, I-VI are given by table 1.1. Moreover, there exists a constant $a_0 = a(0) > 1$ such that the following finite limit exists

$$\lim_{\epsilon_1 \searrow 0} \kappa(\epsilon_1) \cdot \frac{1}{\epsilon_1} a_0^{1/\epsilon_1} > 0 \quad . \tag{1.16}$$

The constant a_0 is given explicitly by (4.9), (4.14.c) evaluated at $\alpha = \epsilon = 0$. For the value of the limit, see (5.10).

The 1-per and 1-hom orbits, viewed as sets, depend continuously on ϵ in the obvious sense. Specifically: crossing line III along a one-dimensional curve from II to IV, two 1-per orbits merge and disappear at a saddle-node (or saddle-saddle) type bifurcation. Crossing I or V, one encounters a blue sky catastrophe.

Since the 1-per orbits bifurcate to different sides from the 1-hom curve $\epsilon_2 = 0$ for $\epsilon_1 < 0$ respectively $\epsilon_1 > 0$, we call this bifurcation a resonant side-switching.



Figure 1.3.a: Resonant side-switching



Figure 1.3.b: Resonant homoclinic doubling

Region	Definition	#1-hom	#1-per
0	$\epsilon_1=0, \epsilon_2=0$	1	0
I	$\epsilon_1 > 0, \epsilon_2 = 0$	1	1
II	$\epsilon_1 > 0 , \ 0 < \epsilon_2 < \kappa(\epsilon_1)$	0	2
III	$\epsilon_1 > 0$, $\epsilon_2 = \kappa(\epsilon_1)$	0	1
IV	$\epsilon_1 \in IR$, $\epsilon_2 > \kappa(\epsilon_1)$	0	0
V	$\epsilon_1 < 0$, $\epsilon_2 = 0$	1	0
VI	$\epsilon_1 \in IR$, $\epsilon_2 < 0$	0	1

 Table 1.1:
 Resonant side-switching; numbers of homoclinic and periodic orbits

Theorem B: Resonant homoclinic doubling.

Let $F = F(\alpha, z)$ of class C^{M+5} , $2 \le M \le \infty$ be a generic two parameter vector field with a twisted resonant homoclinic orbit at $\alpha = 0$. Then resonant homoclinic doubling occurs at $\alpha = 0$ (see figure 1.3.b).

In more detail, let Γ be as in theorem A, but twisted, and let U again denote a sufficiently small tubular neighborhood of its closure clos Γ . Then there exists a C^{M} diffeomorphic reparametrization $\epsilon = \epsilon(\alpha)$ and two functions

$$\epsilon_{2} = \kappa_{\iota}(\epsilon_{1}) \begin{cases} = 0 & \epsilon_{1} \leq 0 \\ for & \\ > 0 & \epsilon_{1} > 0 \\ \end{cases}$$
(1.17)

 $\iota \in \{hom, per\}, of class C^M for \epsilon_1 > 0, such that the numbers of N-periodic (N-per) and of N-homoclinic (N-hom) orbits in U with <math>N = 1, 2$ for the parameter regions 0, I-VIII are given in table 1.2. The curves κ_{ι} have the universal limit property

$$\lim_{\epsilon_1 \searrow 0} \frac{\kappa_{\text{hom}}(\epsilon_1)}{\kappa_{\text{per}}(\epsilon_1)} = \frac{e}{2} = 1.36 \dots \qquad (1.18)$$

Moreover, there exists a constant $a_0 = a(0) < -1$ such that the following finite limit exists

$$\lim_{\epsilon_1 \searrow 0} \kappa_{\text{per}}(\epsilon_1) |a_0|^{1/\epsilon_1} > 0 . \qquad (1.19)$$

The constant a_0 is given explicitly by (4.9), (4.14.c) evaluated at $\alpha = \epsilon = 0$. For the value of the limit, see (6.8).

All homoclinic and periodic orbits depend continuously on ϵ as sets. Specifically: crossing lines I, V, VII yields blue sky catastrophes. Crossing line III, one encounters a period doubling bifurcation of the 1-per orbits to a sheet of 2-per orbits.

The set of 2-per orbits contains a two-dimensional continuum C of orbits which extends from the period doubling 1-per orbits at parameter curve III to the blue sky catastrophe at the curve of 2-hom orbits along line V.

Region	Definition	#1-hom	#2-hom	#1-per	#2-per
0	$\epsilon_1=0, \epsilon_2=0$	1	0	0	0
I	$\epsilon_1 > 0, \epsilon_2 = 0$	1	0	0	0
II	$\epsilon_1 > 0, \ 0 < \epsilon_2 < \kappa_{ m per}(\epsilon_1)$	0	0	1	≥ 0
III	$\epsilon_1 > 0, \epsilon_2 = \kappa_{\mathrm{per}}(\epsilon_1)$	0	1	1	≥ 0
IV	$\epsilon_1 > 0, \ \kappa_{per}(\epsilon_1) < \epsilon_2 < \kappa_{hom}(\epsilon_1)$	0	0	1	≥ 1
V	$\epsilon_1 > 0, \epsilon_2 = \kappa_{\mathrm{hom}}(\epsilon_1)$	0	0	1	≥ 0
VI	$\epsilon_1 \in IR, \epsilon_2 > \kappa_{\mathrm{hom}}(\epsilon_1)$	Ô	0	1	≥ 0
VII	$\epsilon_1 < 0, \epsilon_2 = 0$	1	0	0	0
VIII	$\epsilon_1 \in IR, \epsilon_2 < 0$	0	0	0	0

Table 1.2: Resonant homoclinic doubling; numbers of N-hom and N-per orbits for N = 1, 2.

By a continuum, we mean a relatively compact connected set. For our notion of dimension of C, we refer to [ALEXANDER & ANTMAN] and to lemma 6.5 below.

Our statements concerning 2-per solutions are somewhat weak. Actually, 2-per solutions can occur only in a wedge region which is shaped like the wedge region IV between κ_{per} and κ_{hom} but which is somewhat wider. For a precise statement see lemma 6.4 below. Due to the topological methods which we use, we cannot determine the exact number of 2-per solutions in these regions.

Both theorems, A and B, are new results as they stand. The basic underlying idea, however, that resonant principal eigenvalues $\mu_0 = \nu_0$ can lead to bifurcation, is not new. The non-twisted case was studied before by [LEONTOVICH, NOZDRACHEVA, SANDERS & CUSHMAN] (1951, 1979, 1984) for planar vector fields ($z \in \mathbb{R}^2$). We refer to section 7 for a more detailed discussion.

At this stage, it is high time to mention the pioneering results by Yanagida [YANAGIDA] (1986) on homoclinic doubling in the case of a single unstable dimension (n = 1). In a somewhat less geometric setting Yanagida already gave algebraic conditions for existence and inexistence of 2-hom orbits. In the resonant case, these conditions amount to our twist condition. The proofs, however, are based on local C^1 -linearization in the spirit of the Grobmann-Hartman theorem. But such a linearization may be inappropriate at resonant eigenvalues. Still, we devote section 2 to an exposition of the (non-rigorous) approach by complete

linearization, for illustration purposes. The gap concerning C^1 -linearization was pointed out and closed in \mathbb{R}^3 by Kokubu, see [KOKUBU 1,2] (1987, 1988). These results, although based on "Shilnikov variables", are obtained by a methodology which differs from our approach. For a more detailed discussion we refer to section 7.

Replacing complete linearization, we build our approach on a careful analysis of Shilnikov's parametrization of the flow near the origin; see [SHILNIKOV 5, DENG 1] (1968, 1987). This background material is surveyed in section 3. In section 4, we then derive a Ljapunov-Schmidt type reduction of the bifurcation problem down to an N-dimensional system of bifurcation equations, for N-per and N-hom solutions. These systems are highly transcendental: the bifurcation parameter enters as an exponent in the leading term. This fact causes the exponential smallness of the curves κ in theorems A and B. Theorem A is proved in section 5, discussing the case N = 1. For a proof of theorem B, alias N = 2, see section 6. After a detailed discussion in section 7, we conclude with an appendix which recalls the explicit form of all our genericity assumptions.

2. A model example: the piecewise linear case

In this section, we analyse the flow near a resonant homoclinic orbit Γ in \mathbb{R}^3 . Following Yanagida's idea [YANAGIDA], we assume that our flows are piecewise linear. Due to resonance, this assumption is inadequate to the nonlinear problem. However, deviating from [YANAGIDA], we will treat the piecewise linear case in such a way that it becomes a paradigm to our treatment of the general generic case.

Consider a homoclinic orbit Γ of a three-dimensional vector field $F = F(\alpha, z)$ at $\alpha = 0$, as in theorems A and B. Assume $F(\alpha, 0) = 0$, for all α . We introduce coordinates z = (x, y), $x = x_0$, $y = (y_0, y_1)$ associated to the eigendirections of the eigenvalues $-\mu_0(\alpha) < 0 < \nu_0(\alpha) < \nu_1(\alpha)$ of the linearization $D_z F(\alpha, 0)$. We may rescale time so that $\mu_0(\alpha) = 1$. Moreover, we assume that $\nu_0(\alpha) = 1 + \alpha_1$.



Figure 2.1: Sections and return maps

Our first linearity assumption is the following:

$$F(\alpha, z) = D_z F(\alpha, 0) \cdot z \quad \text{for } |z| = \max(|x|, |y|) \leq \delta_0 \quad . \tag{2.1}$$

In other words, the flow near z = 0 is given by:

$$\dot{x_0} = -x_0 \dot{y_0} = (1 + \alpha_1) y_0 \dot{y_1} = \nu_1 y_1$$
 (2.2)

Let $S_{in} := \{(x, y) : x_0 = \delta_0\}$, $S_{out} := \{(x, y) : y_0 = \delta_0\}$ denote (local) Poincaré sections, transverse to the homoclinic orbit Γ at $\mathbf{q} = (\delta_0, 0, 0)$, $\mathbf{p} = \{0, \delta_0, 0\}$ (cf. Figure 2.1). Note that we assume here that \mathbf{p} lies in the linear eigenspace of ν_0 for simplicity. The linear flow in the box $\{|z| \leq \delta_0\}$ defines a Poincaré map \prod^{loc} from a suitable subset \bar{S}_{in} of S_{in} to S_{out} :

$$\begin{aligned} \prod^{\text{loc}} : S_{\text{in}} &\to S_{\text{out}} \\ (x^{\text{in}}, y^{\text{in}}) &\to (x^{\text{out}}, y^{\text{out}}) . \end{aligned}$$
 (2.3)

Likewise, there is an outer Poincaré map \prod^{far} from S_{out} to S_{in} , given by the flow along Γ :

$$\Pi^{\text{far}}: S_{\text{out}} \to S_{\text{in}}$$

$$(x^{\text{out},j}, y^{\text{out},j}) \to (x^{\text{in},j+1}, y^{\text{in},j+1})$$

$$(2.4)$$

The superscripts j, j + 1 are introduced here because we will be interested in iterates of $\prod^{\text{far}} \circ \prod^{\text{loc}}$ later on.

We also assume that \prod^{far} is (affine) linear,

$$\begin{pmatrix} y_0^{\text{in},j+1} \\ y_1^{\text{in},j+1} \end{pmatrix} = \begin{pmatrix} \alpha_2 \\ 0 \end{pmatrix} + \begin{pmatrix} \Pi^{11} & \Pi^{12} \\ \Pi^{21} & \Pi^{22} \end{pmatrix} \begin{pmatrix} x_0^{\text{out},j} \\ y_1^{\text{out},j} \end{pmatrix}, \text{ with } \Pi^{22} \neq 0 , \quad (2.5)$$

where, again for simplicity, the Π^{ik} are considered now as being independent of α . The generic assumption $\Pi^{22} \neq 0$ guarantees that W^u hits W^s in **q** in general position so that (1.13) holds. Indeed, $T_0 W^{uu}$ is the y_1 -axis. Note the meaning of $\alpha = (\alpha_1, \alpha_2)$. For $\alpha_2 = 0$, the points

$$\mathbf{p} = (x^{out}, \, \delta_0, \, y_1^{out}) := (0, \, \delta_0, \, 0) \in W^u$$

and

$$\mathbf{q} = (\delta_0, y_0^{\text{in}}, y_1^{\text{in}}) := (\delta_0, 0, 0) \in W^2$$

lie on a 1-homoclinic branch, all along the line $\{\alpha_1 \in \mathbb{R}, \alpha_2 = 0\}$. Roughly speaking, α_2 describes the distance between W^u and W^s . The other parameter,

 α_1 , expresses transverse crossing of the eigenvalue $\nu_0 = 1 + \alpha_1$ through $\mu_0 = 1$ along the 1-homoclinic branch $\alpha_2 = 0$.

At this stage, we could proceed by piecing Π^{loc} and Π^{far} together, iterating the return map $\Pi^{\text{far}} \Pi^{\text{loc}}$ of \bar{S}_{in} . Unfortunately, this simple approach does not seem to work equally well in the fully nonlinear case, because, in iterating $\Pi^{\text{far}} \Pi^{\text{loc}}$, it is difficult to keep control of the shrinking cusp-shaped domain of definition in \bar{S}_{in} . In Shilnikov's variables, we obtain nicer domains of definition which are more suitable for applications of the implicit function theorem. Therefore, we now describe the same iterations in Shilnikov's coordinates

$$(s, x^{\text{in}}, y^{\text{out}}) = (s, \delta_0, \delta_0, y_1^{\text{out}})$$
,

[SHILNIKOV 5], rather than using

$$(x^{ ext{in}}, y^{ ext{in}}) = (\delta_0, y^{ ext{in}}_0, y^{ ext{in}}_1) \in S^{ ext{in}}$$
 .

The two coordinate systems are related via the linear flow (2.2) as follows:

$$y_0^{\text{in}} = e^{-\nu_0 s} y_0^{\text{out}} = e^{-\nu_0 s} \delta_0$$

$$y_1^{\text{in}} = e^{-\nu_1 s} y_1^{\text{out}}$$

$$x_0^{\text{out}} = e^{-s} x_0^{\text{in}} = e^{-s} \delta_0 .$$
(2.6)

The parameter s, for Shilnikov time, is the Poincaré time associated to Π^{loc} . For a more detailed exposition of this concept see section 3.

Note that the domain of definition of $\Pi^{\text{far}} \Pi^{\text{loc}}$ takes the form of a rectangle in the Shilnikov coordinates (s, y_1^{out}) , namely $s \ge s^*$ large enough and $|y_1^{\text{out}}| \le \delta_0$.

Reintroducing indices j, j + 1 to keep track of the iterations in the Shilnikov coordinates $(s_j, y_1^{\text{out},j})$, we obtain from (2.5), (2.6):

$$e^{-\nu_0 s_{j+1}} \delta_0 = \alpha_2 + e^{-s_j} \prod^{11} \delta_0 + \prod^{12} y_1^{\text{out},j}$$
(2.7.a)

$$e^{-\nu_1 s_{j+1}} y_1^{\text{out}, j+1} = e^{-s_j} \prod^{21} \delta_0 + \prod^{22} y_1^{\text{out}, j} .$$
 (2.7.b)

We are interested in N-periodic solutions near Γ . With $r_j := e^{-\nu_0 s_j} > 0$ this reads

$$\delta_0 r_{j+1} = \alpha_2 + \delta_0 \prod^{11} r_j^{1/\nu_0} + \prod^{12} y_1^{\text{out}, j} , \qquad (2.8.a)$$

$$r_{j+1}^{\nu_1/\nu_0} y_1^{\text{out},j+1} = \delta_0 \prod^{21} r_j^{1/\nu_0} + \prod^{22} y_1^{\text{out},j} , \qquad (2.8.b)$$

with $j \pmod{N}$. Note that $r_j \delta_0 = y_0^{\text{in},j}$ measures the vertical distances of the periodic trajectory from the stable manifold. Recall that we are interested in solutions with small r_j , $y_1^{\text{out},j}$. In fact a solution with

$$r_0 = 0$$
, and $r_1, \ldots, r_{N-1} > 0$

all distinct, is an N-homoclinic orbit.

Mimicking Ljapunov-Schmidt reduction, we can solve equations (2.8.b) for $(y_1^{\text{out},j}), j = 0, \ldots, N-1$, since \prod^{22} is nonzero by assumption (2.5). We obtain

$$y_1^{\text{out},j} = -\delta_0 \frac{\prod^{21}}{\prod^{22}} r_j^{1/\nu_0} + O(\delta_0 |\mathbf{r}|^{(\nu_1+1)/\nu_0}) \quad , \tag{2.9}$$

where $\mathbf{r} = (r_0, \ldots, r_{N-1})$ and $|\mathbf{r}| := \max r_j$. Plugging this into the remaining equations (2.8.a) we find the bifurcation equations

$$r_{j+1} = \epsilon_2 + a_0 r_j^{1+\epsilon_1} + O\left(|\mathbf{r}|^{(\nu_1+1)/\nu_0}\right) \quad . \tag{2.10}$$

Here we use the notation

$$\begin{array}{rcl} 1/\nu_0 &=& 1/(1+\alpha_1) =: 1+\epsilon_1 \ ,\\ a_0 &=& (\det \ \Pi^{ij})/ \ \Pi^{22} \ ,\\ \epsilon_2 &:=& \alpha_2/\delta_0 \ . \end{array}$$

As a variant, we mention the scaling $\hat{r}_j := e^{-s_j}$. We then obtain analogously

$$\hat{r}_{j} = \hat{\epsilon}_{2} + \hat{a}_{0} \, \hat{r}_{j+1}^{1+\epsilon_{1}} + O(|\hat{\mathbf{r}}|^{\nu_{1}+1}) \quad . \tag{2.10.}$$

where $1 + \hat{\epsilon}_1 := 1 + \alpha_1 = \nu_0$, $\hat{a}_0 = a_0^{-1}$, and $\hat{\epsilon}_2 = -\hat{a}_0 \alpha_2 / \delta_0$.

We are simply amazed at the fact, that virtually the same form (2.10), $(2.10.^{\circ})$ of the iteration emerges from a careful Ljapunov-Schmidt reduction analysis of the fully nonlinear system; see (4.13), $(4.13.^{\circ})$ in corollary 4.3 and remark 4.4.

The nonzero coefficient a_0 in (2.10) has an interesting geometric interpretation. We claim

$$a_0 > 0$$
 if Γ is a non-twisted homoclinic orbit ;
 $a_0 < 0$ if Γ is twisted. (2.11)

This follows directly from our definition 1.1 of a twist. See figures 1.2 and 2.1. Indeed, let e_1 denote the unit vector along the positive y_1 -axis; e^+ , $\dot{\mathbf{p}}$ point along the positive y_0 -axis and e^- , $-\dot{\mathbf{q}}$ along the positive x_0 -axis. Note that the orientations of the triples $(e^-, e_1, \dot{\mathbf{p}})$ and of $(e^+, e_1, \dot{\mathbf{q}})$ are equal:

$$\operatorname{sgn} \left(e^{-} \wedge e_{1} \wedge \dot{\mathbf{p}} \right) = \operatorname{sgn} \left(e^{+} \wedge e_{1} \wedge \dot{\mathbf{q}} \right) = +1 \quad , \tag{2.12}$$

where \wedge denotes the exterior product. Consequently det $\prod^{ij} > 0$, by definition of \prod^{ij} , cf. (2.5). Now note that the orientation of $T_{\mathbf{p}}$, given by $e_1 \wedge \dot{\mathbf{p}}$, continues to that of $T_{\mathbf{q}}$, given by $(\prod^{12} e^+ + \prod^{22} e_1) \wedge \dot{\mathbf{q}}$. Therefore, (2.12) implies

$$\operatorname{sgn} \left(e^{+} \wedge (\Pi^{12} e^{+} + \Pi^{22} e_{1}) \wedge \dot{\mathbf{q}} \right) = \operatorname{sgn} \Pi^{22} \cdot \operatorname{sgn} \left(e^{+} \wedge e_{1} \wedge \dot{\mathbf{q}} \right) =$$

$$= \operatorname{sgn} \Pi^{22} \cdot \operatorname{sgn} \left(e^{-} \wedge e_{1} \wedge \dot{\mathbf{p}} \right) =$$

$$= \operatorname{sgn} a_{0} \cdot \operatorname{sgn} \left(e^{-} \wedge e_{1} \wedge \dot{\mathbf{p}} \right) ,$$

$$\operatorname{since} a_{0} = \left(\det \Pi^{ij} \right) / \Pi^{22} \text{ and } \det \Pi^{ij} > 0 .$$

$$(2.13)$$

Thus e^- and e^+ , respectively, are on the same side of T_p and T_q if $a_0 > 0$. Likewise, they are on opposite sides if $a_0 < 0$. This proves our claim (2.11).

With this new insight, let us now return to the bifurcation equations (2.10), (2.10.[^]). Let us simple-mindedly neglect higher order terms O, altogether. We are only interested in the bifurcation diagrams in parameter space, as given in theorems A, B and figures 1.3.a,b. Therefore, we can assume $|a_0| > 1$ in (2.10). Indeed, $|a_0| = 1$ is nongeneric, and $|a_0| < 1$ is equivalent to $|a_0| > 1$ when switching to (2.10.[^]) and reversing the direction of iteration. It is now fairly straightforward to derive all results of theorems A, B in this simplified situation. All we have to analyse is an iteration of the monotone and convex/linear/concave map

$$r \longrightarrow \epsilon_2 + a_0 r^{1+\epsilon_1}$$

In fact, tables 1.1 and 1.2 then list all possible N-per and N-hom orbits. We leave this simple case to the reader. For a reference see [GLENDINNING]. The analysis for O-terms included fills sections 5 and 6. In particular, we lose control over N-per orbits with $N \ge 2$ in the non-twisted and $N \ge 3$ in the twisted case, due to these error terms. In the twisted case, the number of 2-per orbits can only be estimated. Basically, the reason is that the O-term depends not only on r_j but also on the r_k with $k \ne j$. Thus (2.10) cannot be interpreted as the iteration of a scalar function $r_j \rightarrow r_{j+1}$ anymore. For a more detailed discussion and a remedy see section 7.

3. Shilnikov variables and the strong λ -Lemma

In this section, we briefly review some fundamental (but nontrivial) facts about the Shilnikov variables. These facts are basic to our Ljapunov-Schmidt reduction for resonant bifurcation of homoclinic orbits. In particular, we need careful estimates on derivatives so that we can keep track of higher order error terms during the reduction in section 4. We also give a precise version of the strong λ -Lemma, which was used in our definition of twisted versus non-twisted homoclinic orbits (see Definition 1.1). For a more detailed account, including proofs, we refer to [DENG 1].

We fix the following normalized setting for the rest of this paper. We locally describe the original vector field

$$\dot{z} = F(\alpha, z) \tag{1.1}$$

in suitable coordinates $z = (x, y) \in \mathbb{R}^m \times \mathbb{R}^n$. Specifically (x, y) are chosen such that, near z = 0, takes the form

$$\dot{x} = A(\alpha)x + f(\alpha, x, y) \dot{y} = B(\alpha)y + g(\alpha, x, y) ,$$

$$(3.1)$$

where f, g and their first derivatives $D_{(x,y)}f$, $D_{(x,y)}g$ with respect to x, y vanish identically at x = 0, y = 0, for any (small) $|\alpha|$. Moreover, the linearizations $A(\alpha), B(\alpha)$ are assumed to have block diagonal form

$$A(\alpha) = \left(\begin{array}{c|c} -1 \\ \hline \\ A_1(\alpha) \end{array} \right)$$

$$B(\alpha) = \left(\begin{array}{c|c} \nu_0(\alpha) \\ \hline \\ B_1(\alpha) \end{array} \right)$$
(3.2)

corresponding to the eigenspace decomposition

$$\begin{aligned} x &= (x_0, x_1) \in \mathbb{R} \times \mathbb{R}^{m-1} \\ y &= (y_0, y_1) \in \mathbb{R} \times \mathbb{R}^{n-1} \end{aligned}$$
(3.3)

of x and y. We may further assume that the local (un)stable manifolds W_{loc}^{u} , W_{loc}^{s} are given by

$$W_{loc}^{u} = \{x = 0\}$$

$$W_{loc}^{s} = \{y = 0\} .$$
(3.4)

In other words, we have

$$\begin{array}{rcl} f(\alpha,\,0,\,y) &=& 0 \\ g(\alpha,\,x,\,0) &=& 0 \\ \end{array}, \qquad (3.4')$$

independently of α, x, y near zero. Note that f, g, A, B, ν_0 are of class at least C^{M+4} and the spectra (spec) of A_1, B_1 satisfy

Re spec
$$A_1 < -\bar{\mu}_1 < -1$$

Re spec $B_1 > \bar{\nu}_1 > \nu_0 > 0$, (3.5)

uniformly for small $|\alpha|$ according to our spectral assumption (1.6). Also note that we have normalized the principal stable eigenvalue $-\mu_0(\alpha)$ to become -1, by rescaling time.

With this normalization in mind, we can now reconsider figure 2.1 which, in section 1, served us as an illustration of the flow near the origin and near the homoclinic orbit Γ at $\alpha = 0$. The x_1 -components are omitted, for the sake of simplicity. Also, y_1 now denotes an (n-1)-vector. We pick $\delta_0 > 0$ small enough and choose the local sections $S_{in} = \{x_0 = \delta_0\}$, $S_{out} = \{y_0 = \delta_0\}$ in the δ_0 -box $\{|x| \leq \delta_0, |y| \leq \delta_0\}$ where $|\cdot|$ denotes the max-norm. As before, (x^{in}, y^{in}) and (x^{out}, y^{out}) , respectively, denote elements of S_{in} and S_{out} . As we have indicated in section 2, the Shilnikov variables describing the trajectories for the δ_0 -box are x^{in}, y^{out} , and $s \geq 0$; see e.g. [SHILNIKOV 5]. As before, $s \geq 0$ denotes the time it takes to run inside a δ_1 -box from a point $(x^{in}, \cdot) \in S_{in}$ to a point $(\cdot, y^{out}) \in S_{out}$, whenever this time is defined. With the notation

$$S_{\text{in}}^{x} := \{ x^{\text{in}} = (\delta_{0}, x_{1}^{\text{in}}) : |x_{1}^{\text{in}}| \le \delta_{0} \}, \ S_{\text{out}}^{y} := \{ y^{\text{out}} = (\delta_{0}, y_{1}^{\text{out}}) : |y_{1}^{\text{out}}| \le \delta_{0} \}$$

we have

3.1 Proposition [DENG, Theorems 2.1, 8.1].

Let $\delta_1 > 0$ be chosen small enough, in the above setting. Then there exists a $\delta_0 > 0$ and a unique C^{M+4} -smooth map

$$(x,y) : \{0 < t \le s\} \times S_{in}^{x} \times S_{out}^{y} \times \mathsf{IR}^{2}_{\mathsf{loc}} \to \{(x,y) \in \mathsf{IR}^{m+n} : |x| \le \delta_{1}, |y| \le \delta_{1}\} (3.6)$$
$$(t;s,x^{\mathsf{in}},y^{\mathsf{out}};\alpha) \mapsto (x,y)(t;s,x^{\mathsf{in}},y^{\mathsf{out}};x)$$

such that (x, y), as a function of t, solves the differential equation (3.1) with the boundary conditions

$$\begin{aligned} x &= x^{\text{in}} & at \quad t = 0 \\ y &= y^{\text{out}} & at \quad t = s \quad . \end{aligned}$$
 (3.7)

Smoothness of (x, y) is understood in the sense that (x, y) can be extended smoothly into a neighborhood of the specified time domain, keeping the interpretation of (x, y) being a solution of (3.1). In [DENG 1], a C^k -version was stated, but the C^{∞} -version follows analogously.

Next, we are interested in the limit $s \to \infty$. In fact we can solve the boundary value problem (3.7) for $s = \infty$, at least formally, by letting x, y follow the trajectory of x^{in} in the stable manifold $W_{\text{loc}}^s = \{y = 0\}$ respectively of y^{out} (backwards) in $W_{\text{loc}}^u = \{x = 0\}$. Since we are interested in the quantities $x^{\text{out}}, y^{\text{in}}$, which complement the Shilnikov variables $s, x^{\text{in}}, y^{\text{out}}$ in our δ_0 -box, we now consider

$$\begin{aligned} x^{\text{out}}(r, x^{\text{in}}, y^{\text{out}}; \alpha) &:= x(s; s, x^{\text{in}}, y^{\text{out}}; \alpha) \\ y^{\text{in}}(r, x^{\text{in}}, y^{\text{out}}; \alpha) &:= y(0; s, x^{\text{in}}, y^{\text{out}}; \alpha) , \end{aligned}$$

$$(3.8)$$

where $r := e^{-\nu_0(\alpha)s}$. Note that $r \ge 0$ is Shilnikov time, rescaled so that r = 0 corresponds to $s = \infty$. Similarly, let $\hat{x}^{\text{out}}, \hat{y}^{\text{in}}$ denote the same right-hand sides, but as a function of $\hat{r} := e^{-s}$ instead of r. Then $x^{\text{out}}, y^{\text{in}}, \hat{x}^{\text{out}}, \hat{y}^{\text{in}}$ extend differentiably down to r = 0 as follows.

3.2 Proposition [DENG, Sections 3,4,8].

Let $\delta_0 > 0$ be chosen small enough, in the above setting, and choose any constant ω such that

$$0 < \omega < \min_{\alpha} \{ \nu_0(\alpha), \mu_0(\alpha), \bar{\nu}_1 - \nu_0(\alpha), \bar{\mu}_1 - \mu_0(\alpha) \} ,$$

where the principal eigenvalue $-\mu_0(\alpha)$ was normalized to be -1, above, and $\bar{\mu}_1, \bar{\nu}_1$ bound the remaining (non-principal) eigenvalues as in (1.6).

Then for any finite $0 \le k \le M + 1$ there exits a constant C, which is independent of s, $x^{\text{in}} \in S^x_{\text{in}}, y^{\text{out}} \in S^y_{\text{out}}$, and small α , such that (i)-(iii) below are true.

(i) The following expansion holds for x^{out} as $r \searrow 0$:

$$y^{\text{in}}(r, x^{\text{in}}, y^{\text{out}}; \alpha) = r(\psi(x^{\text{in}}, y^{\text{out}}; \alpha) + R_y)$$
(3.9)

where the remainder term $R_y = R_y(r, x^{\text{in}}, y^{\text{out}}; \alpha)$, of class C^{M+1} for r > 0, is estimated by the C^k -norm $\|\cdot\|_k$ in $(x^{\text{in}}, y^{\text{out}}; \alpha)$:

$$|D_r^{\beta}R_y(r, \cdot, \cdot; \cdot)||_{k-\beta} \le C r^{\omega-\beta} , \qquad (3.10)$$

for $\beta = 0, 1, ..., k$. The leading term $\psi = (\psi_0, \psi_1)$ of class C^{M+1} has the properties:

$$\psi(x^{\rm in}, 0; \alpha) = 0$$
 (3.11.a)

$$\psi_1(0, y^{\text{out}}; \alpha) = 0$$
 (3.11.b)

$$D_{y^{\text{out}}} \psi(0, 0; \alpha) = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}_{n \times n}$$
(3.11.c)

(ii) The corresponding statements for x^{out} , in the same logical context, read

$$x^{\text{out}}(r, x^{\text{in}}, y^{\text{out}}; \alpha) = r^{1/\nu_0(\alpha)}(\varphi(x^{\text{in}}, y^{\text{out}}; \alpha) + R_x) \quad , \tag{3.9}$$

where R_x also satisfies the estimates (3.10). The leading term

 $\varphi = (\varphi_0, \varphi_1) \in C^{M+1}$

satisfies

$$\varphi(0, y^{\text{out}}; \alpha) = 0 \qquad (3.11.a)$$

$$\varphi_1(x^{\rm in}, 0; \alpha) = 0$$
 (3.11.b)

$$D_{x^{\text{in}}}\varphi(0, 0; \alpha) = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}_{m \times m}$$
(3.11.c)

(iii) Replacing $r = e^{-\nu_0(\alpha)s}$ by $\hat{r} = e^{-s}$, that is, replacing $x^{\text{out}}, y^{\text{in}}$ by $\hat{x}^{\text{out}}, \hat{y}^{\text{in}}$, all the above statements remain valid for an expansion

$$\hat{x}^{\text{out}}(\hat{r}, x^{\text{in}}, y^{\text{out}}; \alpha) = \hat{r}(\varphi + \hat{R}_x) ,
\hat{x}^{\text{in}}(\hat{r}, x^{\text{in}}, y^{\text{out}}; \alpha) = \hat{r}^{\nu_0(\alpha)}(\psi + \hat{R}_y) .$$
(3.9)

Occasionally, we refer to (3.9) as Shilnikov's expansion since expansions of this type were first introduced by Shilnikov; see e.g. [SHILNIKOV 5, (2.15)] (1968).

To illustrate the geometric significance of proposition 3.2, we mention that the local strong stable manifold W_{loc}^{ss} is given by those $x^{in} \in S_{in}$ for which

$$\varphi_0(x^{\rm in},0;\alpha) = 0$$
 . (3.12)

See [DENG 1, Corollary 4.3]. Note that $\varphi = 0$ on W_{loc}^{ss} , because $\varphi_1(x^{in}, 0; \alpha) = 0$ by property (3.11.b). So the rapid decay of solutions in W^{ss} , stated in (1.7) above, is reflected in the leading term φ of our exponential expansion (3.9) being zero. Actually, W_{loc}^{ss} is of class C^{M+5} and can be written as the graph of a C^{M+5} function over x_1 ; see e.g. [SHUB, BRUNOVSKÝ & FIEDLER 1]. Here we refer to the regularity C^{M+5} of the vector field, which holds before any normalization of W^u , W^s . Similar statements describe W_{loc}^{uu} , given by $\psi_0(0, y^{out}; \alpha) = 0$.

What are those leading terms φ , ψ , in the completely linear case f = 0, g = 0? We obviously get

$$x(t; s, x^{\text{in}}, y^{\text{out}}; \alpha) = e^{A(\alpha)t} x^{\text{in}}$$

Hence

$$\psi(x^{\mathrm{in}}, y^{\mathrm{out}}; \alpha) = (x_0^{\mathrm{in}}, 0) \in \mathsf{IR}^m$$

projects onto the first (principal) x^{in} -coordinate. In other words, φ and ψ select principal components — also in the nonlinear case. Because φ , ψ are the leading terms in the *r*-expansion (3.9) of x^{out} , y^{in} , this is one more reason for calling the eigenvalues $-\mu_0$, ν_0 and their eigendirections *principal*.

We now discuss the strong λ -Lemma, which enters into definition 1.1 of (non-) twisted homoclinic orbits via the convergence assumption (1.13). We can rewrite (1.13) as

$$\lim_{t \to -\infty} T_{z^{\bullet}(t)} W^{s} = T_{0} W^{ss} \oplus \operatorname{span}(e^{+})$$

$$\lim_{t \to +\infty} T_{z^{\bullet}(t)} W^{u} = \operatorname{span}(e^{-}) \oplus T_{0} W^{uu} ,$$
(3.13)

in the notation of section 1. We give a version of the strong λ -Lemma which shows why the second limit holds, generically. The case of W^s is analogous, reversing time. Consider $\mathbf{q} = z^*(T) \in S_{\text{in}}$, see figures 1.2 and 2.1. The component of $W^u \cap S_{\text{in}}$ which contains \mathbf{q} is a C^{M+4} -manifold of dimension n-1. Perturbing W^u slightly, if necessary, we may assume this manifold to be C^{M+4} -parametrized over y_1 . Here we have further normalized our vector field F at $\alpha = 0$ so that locally

$$W_{loc}^{uu} = \{y_0 = 0, x = 0\} , \text{ and likewise}$$

$$W_{loc}^{ss} = \{x_0 = 0, y = 0\} .$$
(3.14)

We denote the local parametrization of $W^u \cap S_{in}$ near q by the map

$$\begin{aligned} (\xi,\eta): \ y_1 \ \to \ (\xi(y_1), \ \eta(y_1)) \ \in \ S_{\rm in} \ \subseteq \ {\sf IR}^m \ \times \ {\sf IR}^n \\ \eta(y_1) \ = \ (\eta_0(y_1), y_1) \\ (\xi(0), \ \eta(0)) \ = \ {\bf q} \ = \ (\xi(0), 0) \ . \end{aligned}$$

Let \tilde{D}_t^{n-1} , $t \ge T$ denote the image, after time t - T of $im(\xi, \eta) \subseteq S_{in}$ under the flow $F, \alpha = 0$. Note that $z^*(t) \in \tilde{D}_t^{n-1}$. Let D_t^{n-1} denote the component of \tilde{D}_t^{n-1}

in the box B_{δ_0} which contains $z^*(t)$. Finally, we recall the notation $\psi = (\psi_0, \psi_1) = \psi(x^{\text{in}}, y^{\text{out}})$ from proposition 3.2. Here, $\alpha = 0$ is suppressed.

3.3 Proposition (Strong λ -Lemma [DENG 1, Theorem 5.1]).

In the above setting, assume that

$$D_{y_0}\psi(\mathbf{q})\notin T_{\mathbf{q}}W^u . \tag{3.15}$$

Then, as $t \to +\infty$, D_t^{n-1} converges to W_{loc}^{uu} in C^k , for any fixed $0 < k \le M + 1$.

More precisely, there exists a δ -box B_{δ} around z = 0 such that for any $\vartheta > 0$ there exists a $t^* \geq T$ with the following property. For any $t \geq t^*$ the set $B_{\delta} \cap D_t^{n-1}$ is given by the image of a map $y_1 \to (\xi^t, \eta^t)(y_1)$, defined for $|y_1| \leq \delta$ with $\eta_1^t(y_1) = y_1$, and the C^k -norm $\|\cdot\|_k$ satisfies

$$\|(\xi^t, \eta^t)\|_k < \vartheta$$
.

From this proposition, the second limit in (3.13) follows whenever the nondegeneracy condition (3.15) holds. Indeed,

$$T_{z^{\bullet}(t)}W^{u} = \operatorname{span}(\dot{z}^{*}(t)) \oplus T_{z^{\bullet}(t)}D_{t}^{n-1}$$

The first space on the right limits onto $\operatorname{span}(e^{-})$ because $z^{*}(T) = \mathbf{q} \notin W^{ss}$, see (1.11). The second component, by (3.15) and the above proposition, limits onto

$$\{x = 0, y_0 = 0\} = T_0 W^{uu}$$

Incorporating time reversal, both claims of (3.13) are therefore proven.

The nondegeneracy condition (3.15) is a generic condition since we can always enforce (3.15) by a slight perturbation of the vector field F at $\alpha = 0$. Indeed, assume that δ_0 is chosen small enough and leave F unchanged inside the δ_0 -box; this fixes $D_{y_0} \psi(\mathbf{q})$ close to e^+ . But, modifying F slightly along the homoclinic orbit Γ between $\mathbf{p} \in S_{\text{out}}$ and $\mathbf{q} \in S_{\text{in}}$, we can adjust $T_{\mathbf{q}}W^u$ so that it does not contain $D_{y_0}\psi(\mathbf{q})$. Therefore, (3.15) is satisfied generically.

4. Ljapunov - Schmidt reduction

In this section, we derive a reduced normal form equation for N-periodic and N-homoclinic orbits of the vector field $\dot{z} = F(\alpha, z)$ near the homoclinic orbit Γ at $\alpha = (\alpha_1, \alpha_2) = 0$. We describe the flow near the origin z = 0 in Shilnikov's variables $x_1^{\text{in},j}, y_1^{\text{out},j}, r_j = e^{-\nu_0(\alpha)s_j}$, using the setting and the normalizations of section 3. As before, j refers to iterations of the total Poincaré return map

$$\Pi^{\text{tot}} = \Pi^{\text{far}} \circ \Pi^{\text{lo}}$$

where

$$\Pi^{\text{loc}}: S_{\text{in, loc}} \longrightarrow S_{\text{out}}$$

$$(x^{\text{in},j}, y^{\text{in},j}) \longrightarrow (x^{\text{out},j}, y^{\text{out},j}) ,$$

$$\Pi^{\text{far}}: S_{\text{out, loc}} \longrightarrow S_{\text{in}}$$

$$(x^{\text{out},j}, y^{\text{out},j}) \longrightarrow (x^{\text{in},j+1}, y^{\text{in},j+1}) .$$

denote the respective flow-defined maps on suitable domains. For an N-periodic orbit, we have to solve the system

$$\Psi_{j} = \Psi_{j}(\mathbf{r}, \mathbf{x}_{1}^{\text{in}}, \mathbf{y}_{1}^{\text{out}}; \alpha) := \begin{pmatrix} x_{1}^{\text{in}, j+1} \\ y_{0}^{\text{in}}(r_{j+1}, x_{1}^{\text{in}, j+1}, y_{1}^{\text{out}, j+1}; \alpha) \\ y_{1}^{\text{in}}(r_{j+1}, x_{1}^{\text{in}, j+1}, y_{1}^{\text{out}, j+1}; \alpha) \end{pmatrix} - \prod \left(x^{\text{out}}(r_{j}, x_{1}^{\text{in}, j}, y_{1}^{\text{out}, j}; \alpha), y_{1}^{\text{out}, j}; \alpha \right) = 0 \in S_{\text{in}} ,$$

$$(4.1)$$

for all $j(\mod N)$.

Here **r**, $\mathbf{x_1^{in}}$, $\mathbf{y_1^{out}}$, denote the vectors with j-th "component" $r_j > 0$, $x_1^{\text{in},j} \in \mathbb{R}^{m-1}, y_1^{\text{out},j} \in \mathbb{R}^{n-1}$. (If $r_j = 0$ for some j, then we have found a homoclinic orbit.) The map \prod denotes \prod^{far} with the trivial x_0 -component $\prod_{x_0}^{\text{far}} = \delta_0$ omitted. Similarly, x_1^{in} , y_1^{out} can be thought of as being augmented by their trivial components

$$x_0^{\mathrm{in}} = \delta_0$$
 , $y_0^{\mathrm{out}} = \delta_0$,

whenever this is appropriate. We further normalize coordinates so that

$$x_1^{in} = 0$$
 resp. $y_1^{out} = 0$ for
our original homoclinic orbit I
at $\alpha = 0$.

In other words, $(\mathbf{r}, \mathbf{x}_1^{\text{in}}, \mathbf{y}_1^{\text{out}}, \alpha) = 0$ is a trivial solution of $\Psi = 0$, where Ψ has components Ψ_j , of course. We recall that, due to all the above normalizations, both our vector field F and the map Ψ are of class C^{M+4} .

In lemma 4.1 below, we solve the system

$$(\mathbf{id} - \mathbf{P}^{\circ}) \Psi(\mathbf{r}, \mathbf{x}_{1}^{\mathbf{in}}, \mathbf{y}_{1}^{\mathbf{out}}; \alpha) = 0$$

$$(4.2)$$

for $\mathbf{x}_{1}^{in}(\mathbf{r}, \alpha)$, $\mathbf{y}_{1}^{out}(\mathbf{r}, \alpha)$, locally near the trivial solution, by the implicit function theorem, choosing a suitable projection \mathbf{P}° of rank N. We carefully estimate the dependence on \mathbf{r} , since differentiability with respect to r_{i} breaks down when $r_{i} = 0$.

We then modify \mathbf{P}° slightly to become \mathbf{P}^{α} , a rank N projection near \mathbf{P}° which depends on α . In lemma 4.2, we prove a transcendental expansion for the reduced bifurcation equation

$$\boldsymbol{\Phi}(\mathbf{r},\,\alpha) = \mathbf{P}^{\alpha}\,\boldsymbol{\Psi}(\mathbf{r},\,\mathbf{x}_{1}^{\mathrm{in}}(\mathbf{r},\,\alpha),\,\mathbf{y}_{1}^{\mathrm{out}}\left(\mathbf{r},\,\alpha\right)\,;\alpha) = 0 \quad, \tag{4.3}$$

see (4.8). Transforming parameters α to $\epsilon = \epsilon(\alpha)$ suitably, this expansion takes a normal form (4.13) which is (surprisingly) similar to the expansion (2.10) which we have obtained for the linear case. This normal form is the main goal of the present section. We finish the section by relating the sign of a coefficient in the normal form (4.13) to our geometric definition of twisted and non-twisted homoclinic orbits.

Preparing for proofs, we observe that our normalizations imply

$$x^{out}(r, x_1^{in}, y_1^{out}; \alpha) = 0$$

at $r = 0$, (4.4)
 $y^{in}(r, x_1^{in}, y_1^{out}; \alpha) = 0$

for all pertinent x_1^{in} , y_1^{out} , α . Indeed, this follows directly from the definition of the Shilnikov variables since $W_{\text{loc}}^u = \{x = 0\}$, $W_{\text{loc}}^s = \{y = 0\}$. Less directly, we could also invoke proposition 3.2.

For later use, we repeat that the map Ψ is C^{M+4} with respect to \mathbf{x}_1^{in} , $\mathbf{y}_1^{\text{out}}$, α in its domain of definition. As long as $\mathbf{r} > 0$ holds componentwise, Ψ is also C^{M+3} jointly in \mathbf{r} . When some \mathbf{r} -components tend to zero, we remember from proposition 3.2 that the derivatives of Ψ with respect to \mathbf{x}_1^{in} , $\mathbf{y}_1^{\text{out}}$, α behave continuously, up to any finite order $k \leq M+1$. The first derivative of Ψ with respect to \mathbf{x}_1^{in} , $\mathbf{y}_1^{\text{out}}$ is given by

$$D_{(\mathbf{x_{1}^{in}}, \mathbf{y_{1}^{out}})} \Psi_{j} \cdot (\hat{\mathbf{x}_{1}^{in}}, \hat{\mathbf{y}_{1}^{out}}) =$$

$$= -D_{x} \prod D_{x_{1}} x^{out} \cdot \hat{x}_{1}^{in,j} -$$

$$- (D_{x} \prod D_{y_{1}} x^{out} \cdot \hat{y}_{1}^{out,j} + D_{y_{1}} \prod \cdot \hat{y}_{1}^{out,j})$$

$$+ \begin{pmatrix} \hat{x}_{1}^{in,j+1} \\ D_{x_{1}} y^{in} \cdot \hat{x}_{1}^{in,j+1} + D_{y_{1}} y^{in} \cdot \hat{y}_{1}^{out,j+1} \end{pmatrix} =$$

$$= -D_{y_{1}} \prod \hat{y}_{1}^{out,j} + \begin{pmatrix} \hat{x}_{1}^{in,j+1} \\ 0 \end{pmatrix} \text{ (at } \mathbf{r} = 0) .$$

$$(4.5)$$

Indeed, the derivatives of the functions x^{out} and y^{in} vanish identically at $\mathbf{r} = 0$, by (4.4) above.

We can now define the projection \mathbf{P}° for the Ljapunov-Schmidt reduction, cf. (4.2) above. To define the *j*-th component P_{j}° of \mathbf{P}° , we first claim that

$$D_{(\mathbf{x_1^{in}},\mathbf{y_1^{out}})} \Psi_j$$

has maximal rank (corank = 1 in S_{in}) at

$$(\mathbf{r}, \mathbf{x}_1^{in}, \mathbf{y}_1^{out}, \alpha) = 0 ,$$

that is, at the homoclinic orbit Γ . Provided this claim holds true, we can then define

 P_i° : the orthogonal projection onto the one-dimensional

complement in
$$S_{in}$$
 of the image $im \ D_{(\mathbf{x}_1^{in}, \mathbf{y}_1^{out})} \Psi_j$ (4.6)
at $(\mathbf{r}, \mathbf{x}_1^{in}, \mathbf{y}_1^{out}; \alpha) = 0$.

Note that P_j° denotes the same projection for all j, by definition and by our expression (4.5) for $D_{(\mathbf{x_1^{in}}, \mathbf{y_1^{out}})} \Psi_j$. Moreover, rank $P_j^{\circ} = 1$ and rank $\mathbf{P}^{\circ} = N$.

We now show that our claim actually holds, that is, $D_{(\mathbf{x_1^{in}}, \mathbf{y_1^{out}})} \Psi_j$ does possess maximal rank. By (4.5), it is sufficient to show that

$$D_{y_1} \prod_{y_1}$$
 is an isomorphism,

when \prod_{y_1} denotes the y_1 -component of \prod . To identify $D_{y_1} \prod_{y_1}$ as an isomorphism, we first observe that

$$im \ D_{y_1} \prod = T_q W^u \ \cap \ S_{in} \ .$$

This follows because we have normalized $W^u_{\text{loc}} \cap S^{\text{out}}$ to be given by x = 0, $y_0 = \delta_0$. Secondly, motivated by the strong λ -Lemma, we have assumed in (3.13) that

 $\lim_{t\to\infty} T_{z^*(t)} W^u = \operatorname{span}(e^-) \oplus T_0 W^{uu} .$

Since we have normalized W_{loc}^{uu} in (3.14) to be given by

$$W_{\rm loc}^{uu} = \{x = 0, y_0 = 0\}$$
,

and since $\mathbf{q} = z^*(T)$ for some large enough T, this implies that we have an isomorphism

$$im \ D_{y_1} \prod = T_{\mathbf{q}} W^u \cap S_{in} \longrightarrow T_0 W^{uu} = \{x = 0, y_0 = 0\} ,$$

defined by putting x = 0, y = 0 (that is, orthogonal projection). Therefore, $im D_{y_1} \prod_{y_1}$ is indeed an isomorphism, and our definition of P_j^o is justified.

4.1 Lemma: In the above setting, the system

$$(\mathbf{id} - \mathbf{P}^{\circ}) \Psi(\mathbf{r}, \mathbf{x}_{1}^{\mathbf{in}}, \mathbf{y}_{1}^{\mathbf{out}}; \alpha) = 0$$

$$(4.2)$$

has unique solutions x_1^{in} , y_1^{out} , for any given $r \ge 0$, α , near r = 0, $\alpha = 0$. The functions

$$\begin{aligned} \mathbf{x_1^{in}} &= \mathbf{x_1^{in}} \left(\mathbf{r}, \alpha \right) \\ \mathbf{y_1^{out}} &= \mathbf{y_1^{out}} \left(\mathbf{r}, \alpha \right) \end{aligned}$$

are C^{M+4} in (\mathbf{r}, α) for $\mathbf{r} > 0$. They are continuous in \mathbf{r} and C^k in α for $\mathbf{r} \ge 0$ and any finite $k \le M + 1$.

Putting $\mathbf{r} = 0$, denote

$$\begin{aligned} \mathbf{x_1^{in}}(\alpha) &:= \mathbf{x_1^{in}}(0, \alpha) , \\ \mathbf{y_1^{out}}(\alpha) &:= \mathbf{y_1^{out}}(0, \alpha) . \end{aligned}$$

Then the components

$$\begin{array}{rcl} x_1^{\mathrm{in}}(\alpha) &:= & x_1^{\mathrm{in},j}(\alpha) \\ and & y_1^{\mathrm{out}}(\alpha) &:= & y_1^{\mathrm{out},j}(\alpha) \end{array}$$

are the same for all j. Moreover, the following estimates hold for any small $\gamma > 0$ and $|\mathbf{r}| \leq r^0(\gamma)$, $|\alpha| \leq \alpha^0(\gamma)$:

$$\begin{aligned} |\mathbf{x}_{1}^{in}(\mathbf{r}, \alpha) - \mathbf{x}_{1}^{in}(\alpha)| &= C |\mathbf{r}|^{1-\gamma} \\ |\mathbf{y}_{1}^{out}(\mathbf{r}, \alpha) - \mathbf{y}_{1}^{out}(\alpha)| &= C |\mathbf{r}|^{1-\gamma} \end{aligned}$$
(4.7)

Here $C = C(\gamma)$ denotes a large positive constant and $|\mathbf{r}|$ is the max-norm. If we replace the differences of \mathbf{x}_1^{in} , $\mathbf{y}_1^{\text{out}}$ in (4.7) by their derivatives $D_{\alpha}^i D_{\mathbf{r}}^{\beta}$ of total order up to k, then the same estimate holds, except that $|\mathbf{r}|^{1-\gamma}$ gets replaced by $|\mathbf{r}|^{\omega-\gamma-|\beta|}$ where $|\beta| = \beta_0 + \ldots + \beta_{N-1}$ for the multi-index β .

Proof: Existence, uniqueness and regularity of

$$\mathbf{x_1^{in}(r, \alpha), y_1^{out}(r, \alpha)}$$

follow from the implicit function theorem [BERGER, p. 115] with the remarks preceding the lemma. Here we think of Ψ as being extended to all r in a neighborhood of 0, allowing for components r_j to become negative. In fact, we can just extend the maps

$$(x^{\mathrm{out}}, y^{\mathrm{in}}) = (x^{\mathrm{out}}, y^{\mathrm{in}})(r, x^{\mathrm{in}}, y^{\mathrm{out}}; \alpha)$$

to negative values of $r \in \mathbb{R}$ by their leading terms:

$$(x^{\text{out}}, y^{\text{in}}) := \left(-|r|^{1/\nu_0(\alpha)} \varphi(x^{\text{in}}, y^{\text{out}}; \alpha), r \psi(x^{\text{in}}, y^{\text{out}}; \alpha)\right), \text{ for } r < 0$$

This way, $(x^{\text{out}}, y^{\text{in}})$ remain C^{M+1} in $(x^{\text{in}}, y^{\text{out}}; \alpha)$ and all derivatives depend continuously on r. Thus [BERGER, p. 115] applies. For $\nu_0(\alpha) < 1$, the derivatives up to order M in $(x^{\text{in}}, y^{\text{out}}); \alpha)$ depend C^1 on r.

For $\mathbf{r} = 0$ we have $y^{in} = 0$ and $x^{out} = 0$. Hence (4.2) reads

$$\left(\mathrm{id}-P_{j}^{\mathrm{o}}\right)\left(\left(\begin{array}{c}x_{1}^{\mathrm{in},j+1}\\0\end{array}\right)-\prod(0,\,y_{1}^{\mathrm{out},j}\,;\,\alpha)\right)=0\quad\text{for }j(\mathrm{mod}\ N)\ .\tag{4.2.j}$$

Since (4.2.j) defines the same equation, for each j, the components

$$x_1^{\mathrm{in},j}(lpha), \, y_1^{\mathrm{out},j}(lpha)$$

are indeed independent of j.

It remains to prove the estimates (4.7). We first estimate, for small $\mathbf{r} > 0$, the partial derivative

$$|D_{\mathbf{r}} \Psi_j| \leq |D_{r_{j+1}} y^{\mathrm{in}}| + C |D_{r_j} x^{\mathrm{out}}|$$

$$\leq C(1+|r_j|^{-\gamma}) \leq C |\mathbf{r}|^{-\gamma}$$

according to proposition 3.2. Here and below, C denotes possibly different constants. From the implicit function theorem and the above estimate for $D_{\mathbf{r}} \Psi$ we can now conclude

$$\begin{aligned} |\mathbf{x}_{1}^{\mathrm{in}}(\mathbf{r},\,\alpha) - \mathbf{x}_{1}^{\mathrm{in}}(\alpha)| &\leq \int_{0}^{1} |D_{\mathbf{r}}\mathbf{x}_{1}^{\mathrm{in}}(\tau\mathbf{r},\,\alpha)\mathbf{r}| \, d\tau &\leq \\ &\leq C \int_{0}^{1} \tau^{-\gamma} \, d\tau \cdot \, |\mathbf{r}|^{1-\gamma} \leq C \, |\mathbf{r}|^{1-\gamma} \end{aligned}$$

The estimate for y_1^{out} is analogous. The estimates (4.7) for (higher) derivatives of x_1^{in} , y_1^{out} follow from the higher derivative estimates (3.10) on the functions x^{out} and y^{in} . This completes the proof.

We can now define the projection $P(\alpha)$ which enters into our reduced bifurcation equation

$$\mathbf{\Phi}(\mathbf{r},\alpha) = \mathbf{P}^{\alpha} \Psi(\mathbf{r}, \mathbf{x}_{1}^{\text{in}}(\mathbf{r},\alpha), \mathbf{y}_{1}^{\text{out}}(\mathbf{r},\alpha); \alpha) = 0 \quad .$$
(4.3)

We define the *j*-th component P_j^{α} analogously to the projection P_j^{α} which was defined in (4.6) above. We let P_j^{α} denote the orthogonal projection onto the one-dimensional orthogonal complement in S_{in} of the image

$$im D_{(\mathbf{x_1^{in}, y_1^{out}})} \Psi_j$$
 at $\mathbf{r} = 0$, $\mathbf{x_1^{in} = x_1^{in}(\alpha)}$, $\mathbf{x_1^{out} = x_1^{out}(\alpha)}$.

Let $p(\alpha) \in S_{in}$ be a unit vector which spans this complement. Note that $p(\alpha)$ can be chosen to be C^M in α . Like p_j , the projection P_j^{α} does not depend on j. Describing P_j^{α} by the unit vector $p(\alpha) \in S_{in}$, our reduced bifurcation equation becomes the N-dimensional system

$$\Phi_j(\mathbf{r},\alpha) = p(\alpha)^T \Psi_j(\mathbf{r}, \mathbf{x}_1^{\text{in}}(\mathbf{r}, \alpha), \mathbf{y}_1^{\text{out}}(\mathbf{r}, \alpha); \alpha) = 0 \quad , \quad j \pmod{N} \quad . \tag{4.3.j}$$

Note that all x_1 -components $p_{x_1}(\alpha)$ of $p(\alpha)$ are zero, by construction and by the form (4.5) of $D_{(\mathbf{x_1^{in}}, \mathbf{y_1^{out}})} \Psi_j$ at $\mathbf{r} = 0$. Moreover,

$$p(\alpha)^T D_{y_1} \prod(0, y_1^{\text{out}}(\alpha); \alpha) = 0$$
,

by (4.5) and the definition of $p(\alpha)$. Picking δ_0 small enough, we can therefore assume the component $p_y(\alpha)$ to be close to e^+ , that is, the unit vector in the y_0 direction. Indeed, this is true for $\alpha = 0$ by our discussion of $im D_{y_1} \prod$ preceding lemma 4.1. Hence, it is true for all small $|\alpha|$.

We can now state the long-desired transcendental expansion for the reduced bifurcation function Φ_j with respect to r.

4.2 Lemma: The reduced bifurcation function Φ_j given in (4.3.j) above has the expansion

$$\Phi_{j}(\mathbf{r}, \alpha) = -c_{0}(\alpha) + c_{1}(\alpha) r_{j+1} - c_{2}(\alpha) r_{j}^{1/\nu_{0}(\alpha)} + O(|\mathbf{r}|^{1+\omega}) , \ j \pmod{N} ,$$
(4.8)

with

$$c_{0}(\alpha) := p(\alpha)^{T} \prod(0, y_{1}^{\text{out}}(\alpha); \alpha)$$

$$c_{1}(\alpha) := p_{y}(\alpha)^{T} \psi(x^{\text{in}}(\alpha), y^{\text{out}}(\alpha); \alpha) \qquad (4.9)$$

$$c_2(\alpha) := p(\alpha)^T D_{x^{\text{out}}} \prod (0, y_1^{\text{out}}(\alpha); \alpha) \cdot \varphi(x^{\text{in}}(\alpha), y^{\text{out}}(\alpha); \alpha) ,$$

all of class C^{M} . Here

$$x^{\mathrm{in}}(lpha)=(\delta_0,\,x_1^{\mathrm{in}}(lpha))$$
 , $y^{\mathrm{out}}(lpha)=(\delta_0,\,y_1^{\mathrm{out}}(lpha))$,

as usual. The functions φ and ψ were introduced in proposition 3.2. The expansion (4.8) is understood in the following sense. There exists a small $\omega > 0$ such that the remainder term O is estimated, up to a constant factor, by $|\mathbf{r}|^{1+\omega}$. This estimate holds uniformly for $|\alpha| < \alpha_0(\omega)$ including derivatives in α of order up to any finite $k \leq M$.

For $\mathbf{r} > 0$, all terms in the expansion (4.8) are $C^{\mathcal{M}}$. The higher derivatives $D^{i}_{\alpha} D^{\beta}_{\mathbf{r}} \Phi_{j}$, up to total order k, satisfy the analogously differentiated expansion with the error term replaced by

$$O(|\mathbf{r}|^{1+\omega-|\beta|})$$
,

where $|\beta| = \beta_0 + \ldots + \beta_{N-1}$.

Proof: To prove the expansion (4.8), we write (omitting arguments)

$$\Phi_j = p^T \Psi_j = p_y^T y^{\rm in} - p^T \prod$$

and expand the y^{in} -term and the \prod -term, separately, with respect to r. We consider y^{in} first, using Shilnikov's expansion (proposition 3.2) and the estimate (4.7) from lemma 4.1.

$$y^{\text{in}}(r_{j+1}, x_1^{\text{in}, j+1}(\mathbf{r}, \alpha), y_1^{\text{out}, j+1}(\mathbf{r}, \alpha); \alpha) =$$

= $r_{j+1}(\psi(x^{\text{in}, j+1}(\mathbf{r}, \alpha), y^{\text{out}, j+1}(\mathbf{r}, \alpha); \alpha) + R_y) =$ (4.10)
= $\psi(x^{\text{in}}(\alpha), y^{\text{out}}(\alpha); \alpha) r_{j+1} + O(|\mathbf{r}|^{(1+\omega)})$.

This yields the term $c_1(\alpha) r_{j+1}$ in our expansion. Next we expand $p^T \prod \text{at } \mathbf{r} = 0$:

$$p^{T} \Pi = p(\alpha)^{T} \Pi(x^{\text{out}}(r_{j}, x_{1}^{\text{in},j}(\mathbf{r}, \alpha), y_{1}^{\text{out},j}(\mathbf{r}, \alpha); \alpha), y_{1}^{\text{out},j}(\mathbf{r}, \alpha); \alpha) =$$

$$= p(\alpha)^{T} \Pi(x^{\text{out}}(0, \ldots), y_{1}^{\text{out}}(\alpha); \alpha) +$$

$$+ p(\alpha)^{T} D_{(x^{\text{out}}, y_{1}^{\text{out}})} \Pi(x^{\text{out}}(0, \ldots), y_{1}^{\text{out}}(\alpha); \alpha) \cdot$$

$$\cdot \begin{pmatrix} x^{\text{out}}(r_{j}, x_{1}^{\text{in},j}(\mathbf{r}, \alpha), y_{1}^{\text{out},j}(\mathbf{r}, \alpha); \alpha) \\ y_{1}^{\text{out},j}(\mathbf{r}, \alpha) - y_{1}^{\text{out}}(\alpha) \end{pmatrix}$$

$$+ O(|\mathbf{r}|^{1+\omega}) .$$
(4.11)

Indeed, $x^{\text{out}}(0, \ldots) = 0$ and the terms following $D \prod$ will now be estimated to be small of order $O(|\mathbf{r}|^{1-\gamma})$, where $\gamma > 0$ was chosen small in lemma 4.1. Then their squared norms contribute less than $O(|\mathbf{r}|^{1+\omega})$, proving (4.11). Concerning $y_1^{\text{out},j}(\mathbf{r}, \alpha) - y_1^{\text{out}}(\alpha)$ we have an $O(|\mathbf{r}|^{1-\gamma})$ -estimate from lemma 4.1. The term $x^{\text{out}}(r_j, \ldots)$ can be estimated similarly as $y^{\text{in}}(r_{j+1}, \ldots)$ has been estimated in (4.10) above, that is

$$x^{\text{out}}(r_j, x_1^{\text{in},j}(\mathbf{r}, \alpha), y_1^{\text{out},j}(\mathbf{r}, \alpha); \alpha) =$$

= $\varphi(x^{\text{in}}(\alpha)y^{\text{out}}(\alpha); \alpha)r_j^{1/\nu_0(\alpha)} + O(|\mathbf{r}|^{1+\omega})$ (4.12)

since $\nu_0(0) = 1$. With (4.12) at hand, we can now resume our estimate (4.11), using that $p(\alpha)^T D_{y_1} \prod(0, y_1^{\text{out}}(\alpha); \alpha) = 0$ by the properties of the projection

 $p(\alpha)^T$. We get

$$p^{T} \Pi = c_{0}(\alpha) + p(\alpha)^{T} D_{x^{\text{out}}} \prod(0, y_{1}^{\text{out}}(\alpha); \alpha) \cdot \\ \cdot \varphi(x^{\text{in}}(\alpha), y^{\text{out}}(\alpha); \alpha) r_{j}^{1/\nu_{0}(\alpha)} \\ + O(|\mathbf{r}|^{1+\omega}) = \\ = c_{0}(\alpha) + c_{2}(\alpha) r_{j}^{1/\nu_{0}(\alpha)} + O(|\mathbf{r}|^{1+\omega}) .$$

This proves the transcendental expansion (4.8).

The claims concerning differentiability of the expansion follow from the corresponding differentiability statements in lemma 4.1 and proposition 3.2. The proof is therefore complete.

4.3 Corollary: For generic (normalized) vector fields $F = F(\alpha, z)$ of class C^{M+4} , there exists a diffeomorphic local change of parameters $\epsilon = \epsilon(\alpha) \in \mathbb{R}^2$ of class C^M , such that the bifurcation equation (4.8) takes the equivalent normal form

$$r_{j+1} = \epsilon_2 + a(\epsilon) r_j^{1+\epsilon_1} + O(|\mathbf{r}|^{1+\omega}) , \ j \pmod{N}$$

$$(4.13)$$

with $|a(0)| \neq 0, 1$. The error term has the same meaning as in lemma 4.2.

Specifically, the genericity assumptions and the parameter transformation are given as follows. We define

a)
$$\epsilon_1 := \frac{1}{\nu_0(\alpha)} - 1$$

b) $\epsilon_2 := \frac{c_0(\alpha)}{c_1(\alpha)}$ (4.14)
c) $a(\epsilon) := \frac{c_2(\alpha)}{c_1(\alpha)}$,

under the generic assumptions

a)
$$D_{\alpha} c_0(0)$$
 and $D_{\alpha} \nu_0(0)$ are
linearly independent,

b) $c_1(0) > 0$, (4.15) c) $\left| \frac{c_2(\alpha)}{c_1(\alpha)} \right| \neq 0, 1$.

We note explicitly that $\epsilon = 0$ at $\alpha = 0$, since $\nu_0(0) = 1$ and $c_0(0) = 0$.

Proof: We first prove that $\nu_0 = 1$, $c_0 = 0$ at $\alpha = 0$. Then we indicate why the nondegeneracy assumptions (4.15.a-c) are generic. The proof of our claims about the coordinate transformation $\epsilon = \epsilon(\alpha)$ is then an obvious consequence of the standard inverse function theorem, applied to equations (4.14.a,b) in ϵ and α .

Actually, $\nu_0 = 1$ at $\alpha = 0$ since $\nu_0 = \mu_0$, by our resonance assumption (1.9), and since we have normalized μ_0 to be one. Moreover, Γ is a 1-homoclinic orbit at $\alpha = 0$. Therefore, $\mathbf{r} = 0$ is a solution of the reduced bifurcation equation (4.8) at $\alpha = 0$. (We still remember that infinite Shilnikov time s_j corresponds to both a homoclinic orbit and to $r_i = 0$.) Hence $c_0 = 0$ at $\alpha = 0$.

We now address genericity of our assumptions (4.15.a-c): we claim that (4.15.a-c) hold, possibly after a slight perturbation of the original vector field F which does not affect any of the remaining assumptions. Let us compute $D_{\alpha}c_{0}(0)$ first, for $c_{0}(\alpha)$ as defined in (4.9). We obtain

$$D_{\alpha}c_{0}(0) = p(0)^{T} D_{\alpha} \prod(0, 0; \alpha) |_{\alpha=0} , \qquad (4.16)$$

since $y_1(0) = 0$, $\prod(0, 0; 0) = 0$ and since, by definition of p(0), also

$$p(0)^T D_{y_1} \prod = 0$$
 at $(x^{\text{out}}, y_1^{\text{out}}; \alpha) = (0, 0; 0)$.

On the other hand, $\nu_0(\alpha)$ is the principal unstable eigenvalue of the linearization $D_z F(z, \alpha)$ at the equilibrium z = 0. Adjusting both $D_\alpha D_z F$ and $D_\alpha \prod$ slightly, at these points, linear independence of $D_\alpha c_0(0)$ and $D_\alpha \nu_0(0)$ can easily be achieved.

Next we prove $c_1 = p_y^T \psi > 0$. Indeed, we have already picked δ_0 small enough so that $p(\alpha)$ is close to e^+ , the unit vector in the positive y_0 -direction. See the properties of $p(\alpha)$ listed just before lemma 4.2. Therefore, $c_1(0)$ is close to $\psi_0(0, (\delta_0, 0); 0)$, which is close to $\delta_0 > 0$, by property (3.11.c) of ψ . This proves $c_1 > 0$.

Similarly, we can assume $c_2 = p^T D_{x^{\text{out}}} \prod \cdot \varphi \neq 0$ since $\varphi(x^{\text{in}}(\alpha), y^{\text{out}}(\alpha); \alpha)$ points roughly in the x_0 -direction and since $D_{x^{\text{out}}} \prod(0, y_1^{\text{out}}(\alpha); \alpha)$ can be adjusted accordingly. Adjusting $D_{x^{\text{out}}} \prod$ once more, we can finally guarantee

 $|c_2(0)/c_1(0)| \neq 1$,

completing the proof.

4.4 Remark: We note that we may assume |a(0)| > 1, without loss of generality, in our discussion of the normal form (4.13) in section 5. Indeed, use the alternative scaling $\hat{r}_j := e^{-s_j}$ instead of $r_j := e^{-\nu_0(\alpha)s_j}$. Then expansion (4.8) reads

$$\Phi_{j}(\hat{\mathbf{r}}, \alpha) = -c_{0}(\alpha) + c_{1}(\alpha)\hat{r}_{j+1}^{\nu_{0}(\alpha)} - c_{2}(\alpha)\hat{r}_{j} + O(|\hat{\mathbf{r}}|^{1+\omega}), \ j \pmod{N} , \qquad (4.8.)$$
where the $c_{\iota}(\alpha)$ are defined by (4.9), as before. This yields the normal form

$$\hat{r}_{j} = \hat{\epsilon}_{2} + \hat{a}(\hat{\epsilon}) \, \hat{r}_{j+1}^{1+\hat{\epsilon}_{1}} + 0(|\hat{\mathbf{r}}|^{1+\omega}) \,, \, j(\text{mod } N) \,, \qquad (4.13.^{\circ})$$

where $\hat{\epsilon}$, \hat{a} now relate to α , $\nu_0(\alpha)$ by

a)
$$\hat{\epsilon}_1 := \nu_0(\alpha) - 1$$

b) $\hat{\epsilon}_2 := -\frac{c_0(\alpha)}{c_2(\alpha)}$ (4.14.[^])
c) $\hat{a}(\hat{\epsilon}) := \frac{c_1(\alpha)}{c_2(\alpha)} = \frac{1}{a(\epsilon)}$.

This proves our remark about a(0).

As a direct consequence of the normal form (4.13) for resonant homoclinic bifurcation we obtain corollary 4.5.

4.5 Corollary: Along the line $\epsilon_2 = 0$, there exists a 1-homoclinic orbit (N = 1) given by $r_0 = 0$ (primary homoclinic branch). As ϵ_1 passes through zero along this branch, the principal eigenvalues $-\mu_0 = -1$ and $\nu_0 = (1 + \epsilon_1)^{-1}$ cross their resonance transversely. In the transformed parameter ϵ , this reads

$$-\mu_0(\epsilon) + \nu_0(\epsilon) = 0$$

$$D_{\epsilon_1}(-\mu_0(\epsilon) + \nu_0(\epsilon)) \neq 0 , \quad at \quad \epsilon = 0 .$$

$$(4.17)$$

4.6 Lemma: The twist of the homoclinic orbit Γ determines the sign of $a_0 = a(0)$, namely

 $a_0 > 0$ if Γ is non-twisted, $a_0 < 0$ if Γ is twisted.

Proof: Since $a = c_2/c_1$ and since $c_1 > 0$, by corollary 4.3, (4.15.b), we only have to prove that c_2 is negative if Γ is twisted, and positive if Γ is non-twisted.

To relate $c_2 = p^T D_{x^{out}} \prod \cdot \varphi$ to the twist we observe the following, near $\alpha = 0$. First, $p(\alpha)$ is close to e^+ and spans the orthogonal complement to the space $T_{\mathbf{q}} = T_{\mathbf{q}}W^s + T_{\mathbf{q}}W^u$. This follows from the properties of $p(\alpha)$, which are listed just above lemma 4.2, since $T_{\mathbf{q}}$ is spanned by the x-space and $im D_{y_1} \prod$. Second, φ points in the positive x_0 -direction, again by proposition 3.2. In other words, $(\varphi, 0)$ is close to $\delta_0 e^-$. Third, e^- complements the space $T_{\mathbf{p}} = T_{\mathbf{p}} W^s + T_{\mathbf{p}} W^u$, by the strong λ -Lemma. Consequently, $D_{x^{out}} \prod \cdot \varphi$ complements $T_{\mathbf{q}}$, too.

The proof can now be completed as follows. The complement e^- induces an orientation $e_{\mathbf{p}}$ on $T_{\mathbf{p}}$, say, such that $e_{\mathbf{p}} \wedge e^-$ is positive. Here $e_{\mathbf{p}}$ denotes an alternating linear (m+n-1)-form: the volume form on $T_{\mathbf{p}}$. Following the linearized flow along Γ , $e_{\mathbf{p}}$ continues to an orientation $e_{\mathbf{q}}$ on $T_{\mathbf{q}}$ such that $e_{\mathbf{q}} \wedge D \prod e^-$ is also positive. Since $(\varphi, 0)$ is close to e^- , this implies positivity of $e_{\mathbf{q}} \wedge D_{x^{\text{out}}} \prod \varphi$. In other words, e^- and $D_{x^{\text{out}}} \prod \varphi$ point to the same side of $T_{\mathbf{p}}$ and $T_{\mathbf{q}}$, respectively. Since $p(\alpha)$ is orthogonal to $T_{\mathbf{q}}$, we know on the other hand that

$$c_2(\alpha) = p(\alpha)^T \cdot D_{x^{\text{out}}} \prod \varphi$$

is positive resp. negative if $P(\alpha)$ and $D_{x^{\text{out}}} \prod \varphi$ point to the same resp. to opposite sides of $T_{\mathbf{q}}$. Since $p(\alpha)$ is close to e^+ we can combine all this as follows:

$$sign(e_{\mathbf{q}} \wedge e^{+}) = sign(e_{\mathbf{q}} \wedge p(\alpha)) =$$
$$= sign \ c_{2}(\alpha) \cdot sign(e_{\mathbf{q}} \wedge D_{x^{\text{out}}} \prod \varphi) =$$
$$= sign \ c_{2}(\alpha) \cdot sign(e_{\mathbf{p}} \wedge e^{-}) .$$

By definition 1.1, Γ is non-twisted if sign $(e_q \wedge e^+) = \text{sign } (e_p \wedge e^-)$, and twisted otherwise. Since sign $c_2 = \text{sign } a$, this completes the proof.

5. The side-switching bifurcation

In this section, we complete the proof of theorem A on resonant side-switching. We recall from corollary 4.3 that all 1-homoclinic (1-hom) and 1-periodic (1-per) solutions correspond, locally near $\alpha = 0$ and near the homoclinic orbit Γ , to solutions r, ϵ of the reduced bifurcation equation

$$r = \epsilon_2 + a(\epsilon) r^{1+\epsilon_1} + O(r^{1+\omega}) .$$
(5.1)

Here $\epsilon = \epsilon(\alpha) = (\epsilon_1, \epsilon_2)$ are the new normalized parameters, and $r = e^{-\nu(\alpha)s} \ge 0$ describes the exponentially rescaled Shilnikov time associated to the 1-hom resp. the 1-per orbits. Note that we have written r instead of r_j , since we are only interested in 1-per and 1-hom orbits, in this section.

To prove theorem A, we first observe that r, ϵ_1 parametrize the (local) solution set of (5.1). Indeed, a straightforward application of the implicit function theorem enables us to solve (5.1) for ϵ_2 by a C^0 -function

$$\epsilon_2 = \epsilon_2(r, \epsilon_1) \quad , \quad r \ge 0 \quad , \tag{5.2}$$

near $\epsilon = 0$, r = 0. For r > 0, the function ϵ_2 is C^M .

We now consider the surface (5.2) in some more detail. We recall from lemma 4.6 that $a(\epsilon)$ is positive, since Γ is assumed to be non-twisted. By remark 4.4 above, we can further assume a > 1, by a rescaling argument. This implies that

$$\epsilon_2 = \epsilon_2(r, \epsilon_1) \leq 0 \text{ for } r \geq 0, \epsilon_1 \leq 0,$$

as long as $r \leq \rho_0$ is small enough. Indeed,

$$\epsilon_2 = r(1 - a(\epsilon)r^{\epsilon_1} + O(r^{\omega})) \le 0 \tag{5.3}$$

for $\epsilon_1 \leq 0 \leq r \leq \rho_0$ and suitably small ρ_0 , since a(0) > 1. This proves our claims in the left part ($\epsilon_1 \leq 0$) of region IV; see table 1.1 and figure 1.3.a of a resonant side switching for these claims.

Our strategy for completing the proof of theorem A, now, is the following. In lemma 5.2 below we prove that:

$$\epsilon_2 = \epsilon_2(r, \ \epsilon_1) = 0 \tag{5.4}$$

has a unique nontrivial small solution $r = r(\epsilon_1) > 0$, for small $\epsilon_1 > 0$. Of course, there is also the trivial solution r = 0 which describes the 1-hom branch, cf. corollary 4.5. In lemma 5.3 we show that, for fixed ϵ_1 , the C^M -function

$$r \mapsto \epsilon_2 = \epsilon_2(r, \epsilon_1) , r > 0 ,$$
 (5.5)

has zero derivative only along a C^{M} -curve

$$\epsilon_2 = \kappa(\epsilon_1) \ , \ \epsilon_1 > 0 \ ,$$

where the derivative changes sign. We also show that κ is exponentially flat as required in (1.16). By strict monotonicity of $\epsilon_2(\cdot, \epsilon_1)$ off the curve κ , lemma 5.2 and 5.3 then imply all claims concerning multiplicities in the various regions of table 1.1 and figure 1.3.a. Indeed, figure 1.3.a is just the projection of the surface $\epsilon_2 = \epsilon_2(r, \epsilon_1)$ into (ϵ_1, ϵ_2) -space with fold line κ .

We need the following basic lemma about solutions of a transcendental equation.

5.1 Lemma: Let $b = b(\epsilon)$ be a C¹-function such that b(0) > 1. Consider the equation

$$1 = b(\epsilon) r^{\epsilon_1} + O(r^{\omega}) , \quad \epsilon_1 > 0 , \qquad (5.6)$$

with $\omega > 0$. As before, the error term $O(r^{\omega})$ is understood to allow differentiation. Then, for $\epsilon_1 \leq 0$, equation (5.6) cannot have solutions with small $r \geq 0$. For $\epsilon_1 > 0$, equation (5.6) can be solved for $\rho := r^{\epsilon_1}$ as a C^1 -function of ϵ , locally near $\epsilon = (\epsilon_1, \epsilon_2) = 0$, $\rho(0) = 1/b(0)$, by the implicit function theorem. If moreover $\epsilon_2 = O(\epsilon_1)$ for $\epsilon_1 > 0$, then the following limit exists

$$\lim_{\epsilon_1 \searrow 0} r(\epsilon) \cdot b(0)^{1/\epsilon_1} = \exp\left(-D_{\epsilon_1}b(0)/b(0)\right) \quad . \tag{5.7}$$

Proof: Solving (5.6) for $\rho := r^{\epsilon_1}$ is standard if we extend (5.6) to $\epsilon_1 \leq 0$ formally by defining

$$O(r^{\omega}) = O(
ho^{\omega/\epsilon_1}) := 0 \ \ ext{for} \ \ \epsilon_1 \leq 0 \ \ .$$

We obtain

$$\rho(\epsilon) = \frac{1}{b(0)} \left(1 - \frac{D_{\epsilon}b(0)}{b(0)} \epsilon + o(\epsilon) \right) \quad ,$$

for small $|\epsilon|$. If we also assume $\epsilon_2 = o(\epsilon_1)$, then we have

$$r(\epsilon) = \rho(\epsilon)^{1/\epsilon_1} = b(0)^{-1/\epsilon_1} \left(1 - \frac{D_{\epsilon_1} b(0)}{b(0)} \epsilon_1 + o(\epsilon_1) \right)^{1/\epsilon_1}$$

which proves our convergence claim.

We now return to our reduced bifurcation equation.

$$r = \epsilon_2 + a(\epsilon) r^{1+\epsilon_1} + O(r^{1+\omega}) \quad . \tag{5.1}$$

5.2 Lemma: Along the line $\epsilon_2 = 0$, equation (5.1) has the following solution set (locally near $\epsilon = 0$, r = 0):

- (i) the trivial solution r = 0, for all ϵ_1
- (ii) a nontrivial positive solution $r = r_0(\epsilon_1)$, for $\epsilon_1 > 0$, with limiting behavior

$$\lim_{\epsilon \to 0} r_0(\epsilon_1) \cdot a(0)^{1/\epsilon_1} = \exp\left(-D_{\epsilon_1} a(0)/a(0)\right) .$$
 (5.8)

Proof: For $\epsilon_2 = 0$, we divide (5.1) by r;

$$1 = a(\epsilon) r^{\epsilon_1} + O(r^{\omega}) \quad .$$

Then all claims follow from lemma 5.1 with b := a.

It remains to consider fold points of the surface of 1-per solutions

$$r \mapsto \epsilon_2 = \epsilon_2(r, \epsilon_1) \quad , \tag{5.5}$$

projected onto ϵ -space. In other words, we look for (r, ϵ_1) such that

$$D_r \epsilon_2(r, \epsilon_1) = 0$$
.

5.3 Lemma: Locally near $\epsilon = 0$, r = 0, the fold points (r, ϵ_1) of the 1-per surface are given in the form $r = r_*(\epsilon_1)$, $\epsilon_1 > 0$. The function r_* is of class C^M and has the limiting behavior

$$\lim_{\epsilon_1 \searrow 0} r_*(\epsilon_1) a(0)^{1/\epsilon_1} = \exp\left(-1 - D_{\epsilon_1} a(0) / a(0)\right) \quad . \tag{5.9}$$

At $r = r_*(\epsilon_1)$, the derivative $r \mapsto D_r \epsilon_2(r, \epsilon_1)$ changes sign.

The projected fold curve

$$\epsilon_1 \mapsto \epsilon_2 = \kappa(\epsilon_1) := \epsilon_2(r_*(\epsilon_1), \epsilon_1)$$

is also C^{M} , for $\epsilon_{1} > 0$, and has the limiting behavior

$$\lim_{\epsilon_1 \searrow 0} \kappa(\epsilon_1) \cdot \frac{1}{\epsilon_1} a(0)^{1/\epsilon_1} = exp(-1 - D_{\epsilon_1} a(0)/a(0)) \quad . \tag{5.10}$$

Proof: Solving (5.1) for ϵ_2 , by the implicit function theorem, we have obtained

$$r = \epsilon_2(r, \epsilon_1) + a(\epsilon_1, \epsilon_2(r, \epsilon_1))r^{1+\epsilon_1} + O(r^{1+\omega}) \quad . \tag{5.11.a}$$

Recall that $\epsilon_2 = \epsilon_2(r, \epsilon_1)$ is C^M as long as r > 0. Hence the fold condition, $D_r \epsilon_2 = 0$, holds if and only if in addition

$$1 = (1 + \epsilon_1) a(\epsilon_1, \epsilon_2) r^{\epsilon_1} + O(r^{\omega}) , r > 0 , \qquad (5.11.b)$$

with $\epsilon_2 = \epsilon_2(r, \epsilon_1)$. Indeed,

$$1 = D_r \epsilon_2 + (1 + \epsilon_1) a r^{\epsilon_1} + D_{\epsilon_2} a \cdot D_r \epsilon_2 \cdot r^{1 + \epsilon_1} + O(r^{\omega}) , \qquad (5.11.b')$$

and the term $D_{\epsilon_2}a \cdot D_r\epsilon_2 \cdot r^{1+\epsilon_1}$ can be subsumed in the error term $O(r^{\omega})$, for fixed $0 < \omega < 1$. Note that the right-hand side of (5.11.b) is strictly increasing with r. Hence (5.11.b) can have at most one solution $r = r_*(\epsilon_1)$, and $r \to D_r \epsilon_2(r, \epsilon_1)$ changes sign at $r = r_*(\epsilon_1)$.

More specifically, we can apply lemma 5.1 and solve (5.11.b) for $\rho = r^{\epsilon_1}$, $\epsilon_1 > 0$, as a function of ϵ_1 and ϵ_2 near $\epsilon = 0$. Note that

$$r_*(\epsilon_1)^{\epsilon_1} = \rho(\epsilon_1, \epsilon_2(r_*(\epsilon_1), \epsilon_1))$$
(5.12)

and, by standard differentiation of (5.11.b) with respect to ϵ at $\epsilon = 0$,

$$\begin{array}{rcl} (0,\,0) &=& (a\,\rho,\,0) \,+\,\rho\,D_{\epsilon}\,a \,+\,a\,D_{\epsilon}\,\rho \\ &=& (1,\,0) \,+\,\rho\,D_{\epsilon}\,a \,+\,a\,D_{\epsilon}\,\rho \ , \end{array} \tag{5.13}$$

since $a \rho = 1$ at $\epsilon = 0$. In particular, $r_*(\epsilon_1)$ depends differentiably on $\epsilon_1 > 0$, by (5.12), since $D_r \epsilon_2(r_*(\epsilon_1), \epsilon_1) = 0$. In fact, $r_*(\epsilon_1)$ is still C^M for $\epsilon_1 > 0$.

It remains to prove the exponential asymptotics (5.9), (5.10) for $r_*(\epsilon_1)$ and $\kappa(\epsilon_1) = \epsilon_2(r_*(\epsilon_1), \epsilon_1)$. Of course, we would like to utilize lemma 5.1 again. As a first step, we claim

$$\kappa(\epsilon_1) = \epsilon_2(r_*(\epsilon_1), \epsilon_1) = o(\epsilon_1) , \text{ for } \epsilon_1 \searrow 0 .$$
(5.14)

Indeed, κ extends continuously down to $\kappa(0) := 0$ just as $r_*(\epsilon_1) \in [0, r_0(\epsilon_1)]$ does. Moreover, κ is of class C^M for $\epsilon_1 > 0$ with derivative

$$\kappa'(\epsilon_1) = D_{\epsilon_1} \epsilon_2 = -r_*^{1+\epsilon_1} (D_{\epsilon_1} a + D_{\epsilon_2} a \cdot D_{\epsilon_1} \epsilon_2 + a \log r_*) + O(r_*^{1+\omega}) .$$

Here we have used $D_r \epsilon_1(r^*(\epsilon_1), \epsilon_1) = O$, and we have differentiated (5.11.a) to compute $D_{\epsilon_1} \epsilon_2$. Obviously,

$$\lim_{\epsilon_1\searrow 0}\kappa'(\epsilon_1)=0 \;\;,$$

since $r_*(\epsilon_1) \to 0$ as $\epsilon_1 \searrow 0$. Therefore, κ is differentiable down to $\epsilon_1 = 0$ and $\kappa'(0) = 0$. This proves claim (5.14).

Now, lemma 5.1 with $b := (1 + \epsilon_1)a$ applies to the solution $r = r_*$ of (5.11.b). This implies

$$\lim_{\epsilon_1 \searrow 0} r_*(\epsilon_1) a(0)^{1/\epsilon_1} = \exp\left(-1 - D_{\epsilon_1} a(0) / a(0)\right) ,$$

as was claimed in (5.9), since $\epsilon_2 = o(\epsilon_1)$ by (5.14). To prove (5.10) we compute from (5.11.a)

$$\lim_{\epsilon_{1} \searrow 0} \kappa(\epsilon_{1}) \cdot \frac{1}{\epsilon_{1}} a(0)^{1/\epsilon_{1}} = \lim_{\epsilon_{1} \searrow 0} \epsilon_{2}(r_{*}(\epsilon_{1}), \epsilon_{1}) a(0)^{1/\epsilon_{1}} / \epsilon_{1} =$$

$$= \lim_{\epsilon_{1} \searrow 0} r_{*}(\epsilon_{1}) a(0)^{1/\epsilon_{1}} (1 - a(\epsilon) r_{*}(\epsilon_{1})^{\epsilon_{1}} + O(r_{*}^{\omega})) / \epsilon_{1} =$$

$$= \lim_{\epsilon_{1} \searrow 0} r_{*}(\epsilon_{1}) a(0)^{1/\epsilon_{1}} (1 - a(\epsilon) \rho(\epsilon) + O(r_{*}^{\omega})) / \epsilon_{1} =$$

$$= \lim_{\epsilon_{1} \searrow 0} r_{*}(\epsilon_{1}) a(0)^{1/\epsilon_{1}} ((-\rho(0) D_{\epsilon} a(0) - - a(0) D_{\epsilon} \rho(0)) \cdot \epsilon + o(\epsilon)) / \epsilon_{1} =$$

$$= \lim_{\epsilon_{1} \searrow 0} r_{*}(\epsilon_{1}) a(0)^{1/\epsilon_{1}} (\epsilon_{1} + o(\epsilon)) / \epsilon_{1} =$$

$$= \lim_{\epsilon_{1} \searrow 0} r_{*}(\epsilon_{1}) a(0)^{1/\epsilon_{1}} = \exp(-1 - D_{\epsilon_{1}} a(0) / a(0)) .$$

Here we have used (5.12), (5.13) to introduce ρ and to compute $D_{\epsilon} \rho(0)$. The second last equality holds because $\epsilon_2 = o(\epsilon_1)$, by claim (5.14) above. Thus $\kappa(\epsilon_1)$ is exponentially flat at $\epsilon_1 = 0$, and the proofs of lemma 5.3 and of theorem A are complete.

6. The homoclinic doubling bifurcation

In this section, we complete the proof of theorem B on homoclinic doubling. We recall from corollary 4.3 that all N-hom (homoclinic) and N-per (periodic) solutions with N = 1, 2 correspond, locally near $\alpha = 0$ and near the 1-hom Γ , to solutions (r_0, r_1, ϵ) of the reduced bifurcation equation

$$0 = -r_1 + \epsilon_2 + a(\epsilon)r_0^{1+\epsilon_1} + h_0(r, \epsilon)$$

$$0 = -r_0 + \epsilon_2 + a(\epsilon)r_1^{1+\epsilon_1} + h_1(r, \epsilon).$$
(6.1)

Here $\mathbf{r} = (r_0, r_1)$ are the exponentially rescaled Shilnikov times associated to the 1-per $(r_0 = r_1 > 0)$, the 1-hom $(r_0 = r_1 = 0)$, the 2-per $(0 < r_0 \neq r_1 > 0)$ or the 2-hom $(r_0 = 0 < r_1$ or $r_1 = 0 < r_0)$ orbits. As usual, $\epsilon = \epsilon(\alpha) = (\epsilon_1, \epsilon_2)$ are the new normalized parameters. Since Γ is assumed to be twisted, we can assume a(0) < -1; see remark 4.4 and lemma 4.6. The error terms

$$h_j(\mathbf{r}, \epsilon) = O(|\mathbf{r}|^{1+\omega})$$

have the following symmetry at solutions (\mathbf{r}, ϵ) of (6.1):

$$h_0(r_0, r_1, \epsilon) = h_1(r_1, r_0, \epsilon) \quad . \tag{6.2}$$

Indeed, $\mathbf{r} = (r_0, r_1)$ is a solution whenever $\mathbf{\bar{r}} = (r_1, r_0)$ is a solution, because (6.1) was derived from a Poincaré type section of a flow.

Our strategy of proof for theorem B is the following. In lemma 6.1 we show that the 1-per solutions $r_0 = r_1 = r$ form a (local) sheet $\epsilon_2 = \epsilon_2(r, \epsilon_1) > 0$, parametrized over $r \ge 0$ and ϵ_1 . By monotonicity arguments, this sheet projects one-to-one onto the (local) half-space $\{(\epsilon_1, \epsilon_2) \mid \epsilon_2 > 0\}$. The 1-per sheet extends down to the 1-hom branch r = 0, $\epsilon_2 = 0$, which was found in corollary 4.5. In lemma 6.2, we find a unique C^M -curve $\epsilon = \kappa_{per}(\epsilon_1) > 0$, $\epsilon_1 > 0$, where an ordinary period doubling bifurcation occurs on the 1-per sheet. The curve κ_{per} is shown to have the correct exponential asymptotic behavior for $\epsilon_1 \searrow 0$. Lemma 6.3, on the other hand, detects a unique C^M -curve $\epsilon_2 = \kappa_{hom}(\epsilon_1) > 0$, $\epsilon_1 > 0$, where 2-hom orbits occur. We also show the universal scaling property

$$\lim_{\epsilon_1 \searrow 0} \frac{\kappa_{\text{hom}}(\epsilon_1)}{\kappa_{\text{per}}(\epsilon_1)} = \frac{e}{2} \quad . \tag{6.3}$$

In lemma 6.4, we show that there exist C^M -curves $0 < \underline{\kappa}(\epsilon_1) < \overline{\kappa}(\epsilon_1), \epsilon_1 > 0$, such that 2-per solutions do not occur except possibly for pairs (ϵ_1, ϵ_2) with $\underline{\kappa}(\epsilon_1) < \epsilon_2 < \overline{\kappa}(\epsilon_1)$. Our estimating curves $\underline{\kappa}, \overline{\kappa}$ have the same asymptotics as the curves $\frac{1}{2}\kappa_{\text{per}}$, $2\kappa_{\text{hom}}$, but we cannot force them to coincide with these latter curves. Finally, we show in lemma 6.5, that there exists a two-dimensional continuum C of 2-per solutions (\mathbf{r}, ϵ) which connects the 2-hom solutions on the κ_{hom} -curve to the period doubling solutions on the κ_{per} -curve. This 2-per continuum C is confined to the region of ϵ between the two curves $\underline{\kappa}, \overline{\kappa}$. To find C, we use topological methods which are due to [ALEXANDER & ANTMAN]. Therefore, we cannot estimate the precise number of 2-per solutions.

The following three lemmas are much in the spirit of section 5; we assume that the reader is now familiar with the arguments given there. Our first lemma deals with 1-per/1-hom solutions, that is, solutions of

$$r = \epsilon_2 + a(\epsilon) r^{1+\epsilon_1} + O(r^{1+\omega}) \quad . \tag{6.4}$$

Recall that a(0) < -1, in this section.

6.1 Lemma: Locally near $\epsilon = 0$, r = 0 the solutions of (6.4) can be parametrized over (r, ϵ_1) such that

$$\epsilon_2 = \epsilon_2(r, \epsilon_1) \quad . \tag{6.5}$$

Here ϵ_2 is a C^0 -function of $r \ge 0$ and of ϵ_1 ; it is C^1 in $r \ge 0$, $\epsilon_1 > 0$, and C^M for r > 0.

Moreover, $r \mapsto \epsilon_2(r, \epsilon_1)$ is strictly increasing for any fixed ϵ_1 . Therefore, the set of solutions (r, ϵ) projects one-to-one down to those $\epsilon = (\epsilon_1, \epsilon_2)$ with $\epsilon_2 \ge 0$.

Proof: The parametrization (6.5) together with the stated differentiability properties follows directly from the implicit function theorem, applied to equation (6.4). Concerning differentiability, we of course remember the extension of our equations to r < 0 which was discussed in the proof of lemma 4.1. Strict monotonicity of $\epsilon_2(\cdot, \epsilon_1)$ follows by implicit differentiation of (6.4) with respect to r for r > 0. Indeed,

$$1 = (1 + D_{\epsilon_2} a \cdot r^{1+\epsilon_1} + O(r^{1+\omega})) D_r \epsilon_2 + + (1+\epsilon_1) a(\epsilon) r^{\epsilon_1} + O(r^{\omega})$$

implies that $D_r \epsilon_2$ is positive, for all sufficiently small r, ϵ_1 , since a(0) < 0. This completes the proof.

Next, we are interested in period doubling bifurcations on the 1-per sheet $\epsilon_2 = \epsilon_2(r, \epsilon_1)$. These bifurcations can be detected in system (6.1). The 1-per sheet are those solutions of (6.1) for which $r_0 = r_1 = r$. For r > 0, let

$$\Delta = \Delta(r, \epsilon_1) \tag{6.6}$$

denote the determinant of the 2 × 2-Jacobian of the right-hand side of (6.1) with respect to (r_0, r_1) at $r_0 = r_1 = r$, $\epsilon_2 = \epsilon_2(r, \epsilon_1)$.

6.2 Lemma: The determinant Δ on the 1-per sheet $\epsilon_2 = \epsilon_2(r, \epsilon_1), r > 0$, vanishes only along a unique C^M -curve

$$r = \tilde{r}(\epsilon_1) > 0$$
 , $\epsilon_1 > 0$.

and Δ changes sign there.

Viewed in the projection onto ϵ -space, this curve is given by a C^{M} -curve

 $\epsilon_2 = \kappa_{
m per}(\epsilon_1) := \epsilon_2(ilde{r}(\epsilon_1), \epsilon_1) \ , \ \epsilon_1 > 0 \ .$

The asymptotics of \tilde{r} and κ_{per} for $\epsilon_1 \searrow 0$ are

$$\lim_{\epsilon_1 \searrow 0} \tilde{r}(\epsilon_1) \ |a(0)|^{1/\epsilon_1} = \exp(-1 - D_{\epsilon_1} a(0)/a(0)) \ , \tag{6.7}$$

$$\lim_{\epsilon_1 \searrow 0} \kappa_{\text{per}}(\epsilon_1) \ |a(0)|^{1/\epsilon_1} = 2 \exp(-1 - D_{\epsilon_1} a(0)/a(0)) \ . \tag{6.8}$$

Proof: Differentiating (6.1) with respect to (r_0, r_1) at $r_0 = r_1 = r > 0$ we see that

$$\Delta = \det \left(\begin{array}{cc} (1+\epsilon_1) \, a \, r^{\epsilon_1} & -1 \\ -1 & (1+\epsilon_1) \, a \, r^{\epsilon_1} \end{array} \right) + O(r^{\omega}) \quad .$$

Thus Δ vanishes if and only if

a)
$$1 = (1 + \epsilon_1) a(\epsilon) r^{\epsilon_1} + O(r^{\omega}), \text{ or}$$

b)
$$-1 = (1 + \epsilon_1) a(\epsilon) r^{\epsilon_1} + O(r^{\omega}).$$
(6.9)

Since a < 0, the first equation (6.9.a) cannot have solutions with small r. Solving (6.9.b) together with the original fixed point equation

$$r = \epsilon_2 + a(\epsilon) r^{1+\epsilon_1} + O(r^{1+\omega}) , \qquad (6.4)$$

this time with a(0) < -1, is quite similar to the situation which came up for the fold curve κ in the proof of lemma 5.3; see (5.11.a,b). We only indicate the necessary ramifications.

First, we note that (6.9.b) can have a solution for small r and $|\epsilon_1|$ only when $\epsilon_1 > 0$, since a(0) < -1. As usual, we can then solve (6.9.b) uniquely for $r = \tilde{r}(\epsilon_1)$ at $\epsilon_2 = \epsilon_2(r, \epsilon_1)$, by monotonicity. Likewise, we obtain a differentiable $\rho = \rho(\epsilon_1, \epsilon_2)$ such that, analogously to (5.12),

$$\tilde{r}(\epsilon_1)^{\epsilon_1} = \rho(\epsilon_1, \, \epsilon_2(\tilde{r}(\epsilon_1), \, \epsilon_1)) \quad . \tag{6.10}$$

The function $\tilde{r}(\epsilon_1)$ is of class C^M for $\epsilon_1 > 0$. At $\epsilon = 0$, we have $\tilde{r} = 0$ and

$$(1,0) = \rho D_{\epsilon} a + a D_{\epsilon} \rho , \qquad (6.11)$$

since $a \rho = -1$, this time, in (5.13).

As in (5.14), we claim

$$\kappa_{\text{per}}(\epsilon_1) = \epsilon_2(\tilde{r}(\epsilon_1), \epsilon_1) = o(\epsilon_1) , \text{ for } \epsilon_1 \searrow 0 .$$
(6.12)

Unfortunately, we cannot use $D_r \epsilon_2(\tilde{r}, \epsilon_1) = 0$, this time, since we are not at a fold point. However, to show (6.12), we can first differentiate (6.10) with respect to ϵ_1 to obtain

$$D_{\epsilon_1} \tilde{r}(\epsilon_1) = O\left(\frac{1}{\epsilon_1^2} |\rho|^{1/\epsilon_1}\right) \quad , \quad |\rho| < 1 \quad . \tag{6.13}$$

Here we have used

$$\lim_{\epsilon_1 \searrow 0} D_r \, \epsilon_2(\tilde{r}(\epsilon_1), \, \epsilon_1) = 2 \quad , \tag{6.14}$$

which follows by differentiating (6.4). Also, we have used that

$$\lim_{\epsilon_1\searrow 0} D_{\epsilon_1} \epsilon_2(\tilde{r}(\epsilon_1), \epsilon_1) = 0 ,$$

holds, by the same reasoning as in the proof of (5.14) above. Now we can conclude

$$\lim_{\epsilon_1\searrow 0} \kappa'_{\rm per}(\epsilon_1) = \lim_{\epsilon_1\searrow 0} \left(D_r \,\epsilon_2 \,\cdot\, D_{\epsilon_1} \,\tilde{r} \,+\, D_{\epsilon_1} \,\epsilon_2 \right) = 0 \ ,$$

since the first term was estimated in (6.13-14) above and the second term yields zero, as before. This proves (6.12).

With these preparations, lemma 5.1 applies to (6.9.b) with

$$b(\epsilon) := (1 + \epsilon_1) \cdot (-a(\epsilon))$$

and yields

$$\lim_{\epsilon_1 \searrow 0} \tilde{r}(\epsilon_1) |a(0)|^{1/\epsilon_1} = \exp(-1 - D_{\epsilon_1} a(0)/a(0))$$

This proves expansion (6.7) for $\tilde{r}(\epsilon_1)$.

To prove expansion (6.8) for $\kappa_{per}(\epsilon_1)$, we compute similarly to the proof of lemma 5.3

$$\lim_{\epsilon_{1} \searrow 0} \kappa_{\text{per}}(\epsilon_{1}) |a(0)|^{1/\epsilon_{1}} = \lim_{\epsilon_{1} \searrow 0} \epsilon_{2}(\tilde{r}(\epsilon_{1}), \epsilon_{1}) |a(0)|^{1/\epsilon_{1}} =$$

$$= \lim_{\epsilon_{1} \searrow 0} \tilde{r}(\epsilon_{1}) |a(0)|^{1/\epsilon_{1}} (1 - a(\epsilon) \tilde{r}(\epsilon_{1})^{\epsilon_{1}} + O(\tilde{r}^{\omega})) =$$

$$= \lim_{\epsilon_{1} \searrow 0} \tilde{r}(\epsilon_{1}) |a(0)|^{1/\epsilon_{1}} (1 - a(\epsilon) \rho(\epsilon) + O(\tilde{r}^{\omega})) =$$

$$= 2 \exp (-1 - D_{\epsilon_{1}} a(0)/a(0)) ,$$

since

$$\lim_{\epsilon_1 \searrow 0} a(\epsilon) \rho(\epsilon) = -1 \text{ for}$$

$$\epsilon = (\epsilon_1, \epsilon_2(\tilde{r}(\epsilon_1), \epsilon_1)), \ \rho = \tilde{r}^{\epsilon_1}, \text{ by } (6.9.b)^{+}.$$

Therefore (6.8) holds, and the lemma is proved.

Note that lemma 6.2 implies a (local) period doubling bifurcation from the 1-per sheet. Indeed, fix $\epsilon_1 > 0$ and increase the remaining parameter ϵ_2 through the bifurcation point $\epsilon_2 = \kappa_{per}(\epsilon_1)$. Then the determinant Δ of the linearization of system (6.1) at the corresponding 1-per solutions $r_0 = r_1 = r$ changes sign. This change of topological degree implies local bifurcation of 2-per solutions: a period doubling bifurcation. We will return to these arguments in much more detail in lemma 6.5.

We can also look at the other limiting case of 2-per solutions, namely, at 2-hom orbits. Recall that $r_0 = 0 < r_1$, for 2-hom. (Alternatively, we could of course consider the equivalent case $r_0 > 0 = r_1$.)

6.3 Lemma: 2-hom solutions $r_0 = 0 < r_1$ occur only along a unique C^M -curve

$$\epsilon_2 = \kappa_{
m hom}(\epsilon_1) > 0 \ , \ r_1 = \hat{r}(\epsilon_1) > 0 \ , \ \epsilon_1 > 0 \ .$$

The asymptotics of \hat{r} and κ_{hom} for $\epsilon_1 \searrow 0$ are

$$\lim_{\epsilon_1 \searrow 0} \hat{r}(\epsilon_1) \ |a(0)|^{1/\epsilon_1} = \ exp(-D_{\epsilon_1} \ a(0)/a(0)) \tag{6.15}$$

$$\lim_{\epsilon_1 \searrow 0} \kappa_{\text{hom}}(\epsilon_1) \ |a(0)|^{1/\epsilon_1} = \ \exp(-D_{\epsilon_1} \ a(0)/a(0)) \ . \tag{6.16}$$

In particular, we have the universal limit

$$\lim_{\epsilon_1 \searrow 0} \frac{\kappa_{\text{hom}}(\epsilon_1)}{\kappa_{\text{per}}(\epsilon_1)} = \frac{e}{2} = 1.36\dots$$
(6.17)

Proof: Equations (6.1) with $r_0 = 0$ yield

a)
$$0 = -r_1 + \epsilon_2 + O(r_1^{1+\omega})$$

b) $0 = \epsilon_2 + a(\epsilon)r_1^{1+\epsilon_1} + O(r_1^{1+\omega})$, (6.18)

since $|\mathbf{r}| = r_1$. From (6.18.a) we find

$$\epsilon_2 = \epsilon_2(r_1, \epsilon_1) = r_1 + O(r_1^{1+\omega}) ,$$
 (6.19)

where $O(r_1^{1+\omega})$ depends only on r_1 and ϵ_1 . Plugging this into (6.18.b) and using differentiability of $a(\epsilon)$, we obtain

$$0 = r_1 + a(\epsilon_1, 0) r_1^{1+\epsilon_1} + O(r_1^{1+\omega})$$

or, equivalently,

$$1 = -a(\epsilon_1, 0) r_1^{\epsilon_1} + O(r_1^{\omega}) . \qquad (6.20)$$

Solving (6.20) for $r_1 = \hat{r}_1(\epsilon_1)$, by lemma 5.1 again, we find the asymptotics (6.15) for $\hat{r}(\epsilon_1)$. In particular,

$$\lim_{\epsilon_1\searrow 0}\hat{r}(\epsilon_1)=0$$

Now (6.19) yields the same asymptotics for

$$\kappa_{\rm hom}(\epsilon_1) = \epsilon_2(\hat{r}(\epsilon_1), \epsilon_1)$$

since

$$\lim_{\epsilon_1\searrow 0} \epsilon_2(\hat{r}(\epsilon_1), \epsilon_1) |a(0)|^{1/\epsilon_1} = \lim_{\epsilon_1\searrow 0} \hat{r}(\epsilon_1) |a(0)|^{1/\epsilon_1}$$

This proves the κ_{hom} -asymtotics (6.16). Finally, the limit

$$\lim_{\epsilon_1 \searrow 0} \frac{\kappa_{\text{hom}}(\epsilon_1)}{\kappa_{\text{per}}(\epsilon_1)} = \lim_{\epsilon_1 \searrow 0} \frac{\kappa_{\text{hom}}(\epsilon_1) \cdot |a(0)|^{1/\epsilon_1}}{\kappa_{\text{per}}(\epsilon_1) \cdot |a(0)|^{1/\epsilon_1}} = \frac{e}{2}$$

is universal, by the asymptotics (6.8) and (6.16) for κ_{per} and κ_{hom} . This proves the lemma.

Locally near r = 0, $\epsilon = 0$, the 2-per solutions are confined to an exponentially thin wedge in parameter space. More specifically, we have

6.4 Lemma: Consider system (6.1) in a suitably small, fixed neighborhood of $r = 0, \epsilon = 0$.

Then there exist two C^{M} -curves

$$\epsilon_2 = \underline{\kappa}(\epsilon_1)$$
 and $\epsilon_2 = \overline{\kappa}\epsilon_1, \ \epsilon_1 > 0$

with $\underline{\kappa} < \bar{\kappa}$, such that (6.1) does not possess 2-per or 2-hom solutions for ϵ outside of the wedge

$$\{(\epsilon_1, \epsilon_2) \mid \underline{\kappa}(\epsilon_1) < \epsilon_2 < \overline{\kappa}(\epsilon_1)\}$$
(6.21)

in between $\underline{\kappa}$ and $\overline{\kappa}$. The curves $\underline{\kappa}$ and $\overline{\kappa}$, respectively, have the same exponential asymptotics as the curves $\frac{1}{2}\kappa_{per}$ and $2\kappa_{hom}$ (see lemmata 6.2, 6.3, (6.8) and (6.16)). **Proof:** Suppose system (6.1) does possess a 2-per/2-hom solution $\mathbf{r} = (r_0, r_1)$, that is, $r_0 \neq r_1$ and

$$0 = -r_1 + \epsilon_2 + a(\epsilon)r_0^{1+\epsilon_1} + h_0(r_0, r_1, \epsilon)$$

$$0 = -r_0 + \epsilon_2 + a(\epsilon)r_1^{1+\epsilon_1} + h_0(r_1, r_0, \epsilon).$$
(6.1)

Here we have used that, by the symmetry $r_0 \leftrightarrow r_1$, we know

$$h_1(r_0, r_1, \epsilon) = h_0(r_1, r_0, \epsilon)$$
,

at any solution (r_0, r_1) . We may assume

$$0 \leq r_0 < r_1$$
,

so that both error terms h_0, h_1 are of the order $O(r_1^{1+\omega})$. Taking the difference of the two equations (6.1) and dividing by $(r_1 - r_0) > 0$ we obtain an equation of the form

$$-1 = a(\epsilon) \sigma_{\epsilon_1}(r_0, r_1) + O(r_1^{\omega}) , \qquad (6.22)$$

where σ_{ϵ_1} denotes the slope

$$\sigma_{\epsilon_1}(r_0, r_1) := \frac{r_1^{1+\epsilon_1} - r_0^{1+\epsilon_1}}{r_1 - r_0}$$

of the secant to the graph $r \mapsto r^{1+\epsilon_1}$ at the two points $r = r_0, r_1$. Note that this graph is convex or linear or concave, depending on the sign of ϵ_1 . Thus σ_{ϵ_1} always lies in the interval with endpoints

$$r_1^{\epsilon_1}$$
 and $(1+\epsilon_1)r_1^{\epsilon_1}$, (6.23)

for $0 \le r_0 < r_1$. The error term $O(r_1^{\omega})$ is justified, since

$$h_0(r_1, r_0, \epsilon) - h_0(r_0, r_1, \epsilon) =$$

= $(r_1 - r_0) \int_0^1 (D_1 h_0(r_0 + \Theta(r_1 - r_0), r_1 - \Theta(r_1 - r_0), \epsilon) - D_2 h_0(r_0 + \Theta(r_1 - r_0), r_1 - \Theta(r_1 - r_0), \epsilon)) d\Theta$

where the integral is $O(r_1^{\omega})$, by differentiation of $O(|\mathbf{r}|^{1+\omega})$.

It remains to construct $\underline{\kappa}(\epsilon_1)$, $\overline{\kappa}(\epsilon_1)$ such that (6.22) cannot hold outside the specified wedge (6.21). Obviously, (6.22) cannot hold for $\epsilon_1 \leq 0$, by estimate (6.23) for σ and because a(0) < -1. For $\epsilon_1 > 0$, we can estimate ϵ_2 from the first equation in (6.1) as

$$(1+O(r_1^{\omega}))r_1 \leq \epsilon_2 \leq r_1(1-a(\epsilon)r_1^{\epsilon_1}+O(r_1^{\omega}))$$

and, since $a(\epsilon_1, \epsilon_2)$ is Lipschitz in ϵ_2 , this implies an estimate

$$(1+O(r_1^{\omega}))r_1 \leq \epsilon_2 \leq r_1(1-a(\epsilon_1,0)r_1^{\epsilon_1}+O(r_1^{\omega})) .$$
 (6.24)

Plugging (6.24) into (6.22) yields similarly

$$-1 = a(\epsilon_1, 0) \sigma_{\epsilon_1}(r_0, r_1) + O(r_1^{\omega}) .$$

Let $\underline{C} \leq \mathbf{0} \leq \overline{C}$ denote real constants such that

$$\underline{C} r_1^{\omega} < O(r_1^{\omega}) < \overline{C} r_1^{\omega} ,$$

above and in (6.24). Then

$$(1+\epsilon_1) a(\epsilon_1, 0) r_1^{\epsilon_1} + \underline{C} r_1^{\omega} < -1 < a(\epsilon_1, 0) r_1^{\epsilon_1} + \overline{C} r_1^{\omega} ,$$

where the terms to the right and left of -1 are strictly decreasing with r_1 . Therefore, we conclude

$$r_1 \in (\underline{r}_1, \overline{r}_1)$$

where

$$-1 = a(\epsilon_1, 0) \bar{r}_1^{\epsilon_1} + \bar{C} \bar{r}_1^{\omega} \text{ and}$$

$$-1 = (1 + \epsilon_1) a(\epsilon_1, 0) \underline{r}_1^{\epsilon_1} + \underline{C} \underline{r}_1^{\omega} .$$

This defines $\underline{r}_1(\epsilon_1)$, $\overline{r}_1(\epsilon_1)$. Note that $\underline{r}_1(\epsilon_1)$ has the same exponential asymptotics (6.7) as $\tilde{r}(\epsilon_1)$, which is associated to period doubling. Similarly, $\overline{r}_1(\epsilon_1)$ has the "2-hom" asymptotics (6.15) of $\hat{r}(\epsilon_1)$. These facts follow from lemma 5.1. Now we define

$$\underline{\kappa}(\epsilon_1) := \underline{r}_1(\epsilon_1)(1 + \underline{C}\,\overline{r}(\epsilon_1)^{\omega_1})$$

$$\bar{\kappa}(\epsilon_1) := \overline{r}_1(\epsilon_1)(1 - a(\epsilon_1, 0)\,\overline{r}_1(\epsilon_1)^{\epsilon_1} + \overline{C}\,\overline{r}(\epsilon_1)^{\omega}) .$$

Then (6.24) implies that ϵ_2 lies between $\underline{\kappa}$ and $\overline{\kappa}$:

$$\underline{\kappa}(\epsilon_1) < \epsilon_2 < \overline{\kappa}(\epsilon_1)$$
 .

Moreover, $\underline{\kappa}$ has the same exponential asymptotics as $\frac{1}{2} \kappa_{per}$. Indeed,

$$\lim_{\epsilon_1 \searrow 0} \underline{\kappa}(\epsilon_1) |a(0)|^{1/\epsilon_1} = \lim_{\epsilon_1 \searrow 0} \underline{r}_1(\epsilon_1) \cdot |a(0)|^{1/\epsilon_1} =$$
$$= \exp\left(-1 - D_{\epsilon_1} a(0)/a(0)\right)$$
$$= \lim_{\epsilon_1 \searrow 0} \frac{1}{2} \kappa_{\mathrm{per}}(\epsilon_1) |a(0)|^{1/\epsilon_1} ,$$

by lemma 6.2. Likewise,

$$\lim_{\epsilon_1 \searrow 0} \bar{\kappa}(\epsilon_1) |a(0)|^{1/\epsilon_1} =$$

$$= \lim_{\epsilon_1 \searrow 0} \bar{r}(\epsilon_1) |a(0)|^{1/\epsilon_1} \cdot (1 - a(\epsilon_1, 0) \bar{r}_1(\epsilon_1)^{\epsilon_1} + \bar{C} \bar{r}(\epsilon_1)^{\omega})$$

$$= 2 \exp\left(-D_{\epsilon_1} a(0)/a(0)\right) =$$

$$= \lim_{\epsilon_1 \searrow 0} 2 \kappa_{\text{hom}}(\epsilon_1) |a(0)|^{1/\epsilon_1} ,$$

by lemma 6.3. This proves the lemma.

To complete the proof of theorem B, it remains to detect a 2-dimensional continuum C of 2-per solutions which connects the curve of period doubling solutions to the curve of 2-hom orbits. We construct this continuum using a topological multiparameter result due to [ALEXANDER & ANTMAN].

We briefly describe the basic idea, fixing the setting along our way. Fix small enough bounds $\epsilon_2^0 > 0$ and ρ^0 . Then choose $\epsilon_1^0 > 0$ small enough, depending on these bounds, as detailed below. We consider the box

$$\mathcal{B} = \{ (\epsilon, \mathbf{r}) \mid 0 < \epsilon_2 < \epsilon_2^0, 0 < \epsilon_1 < \epsilon_1^0, 0 < \mathbf{r} < \rho^0 \} ,$$

where the **r**-inequality is understood componentwise, for r_0 and r_1 . Fix ϵ_1 , for a moment. The corresponding section \mathcal{B}_{ϵ_1} of the box \mathcal{B} is drawn in figure 6.1. We see the 1-per branch in the $r_0 = r_1$ diagonal, the period-doubling point $\epsilon_2 = \kappa_{per}(\epsilon_1), r_0 = r_1 = \tilde{r}(\epsilon_1)$ on that 1-per branch, and the two 2-hom points given by $\epsilon_2 = \kappa_{\text{hom}}(\epsilon_1)$ and $r_0 = 0$, $r_1 = \hat{r}(\epsilon_1)$ or $r_0 = \hat{r}(\epsilon_1)$, $r_1 = 0$. By lemma 6.2, the sign of the determinant Δ of the r-linearization of (6.1) changes at $\epsilon_2 = \kappa_{per}$ along the 1-per branch. By the classical global Rabinowitz theorem [RABINOWITZ], this implies the existence of a global bifurcating continuum \mathcal{C}_{ϵ_1} of 2-per solutions $(r_0 \neq r_1)$ from the 1-per branch. Since $\epsilon_2 = \kappa_{per}$ is the only bifurcation point, this continuum \mathcal{C}_{ϵ_1} extends to the boundary of the open box \mathcal{B}_{ϵ_1} . Since \mathcal{C}_{ϵ_1} consists of 2-per solutions, the values of ϵ_2 on \mathcal{C}_{ϵ_1} are confined to the wedge region $\underline{\kappa}(\epsilon_1) < \epsilon_2 < \bar{\kappa}(\epsilon_1)$. Of course, we assume here that ϵ_1^0 is chosen such that $\bar{\kappa}(\epsilon_1) < \epsilon_2^0$ for $0 < \epsilon_1 < \epsilon_1^0$. Since C_{ϵ_1} extends to the boundary bdy \mathcal{B}_{ϵ_1} , the closure clos \mathcal{C}_{ϵ_1} , intersected with bdy \mathcal{B}_{ϵ_1} , consists precisely of the two 2-hom points. The following lemma gives a 2-dimensional account of this 1-dimensional result, including the parameter ϵ_1 which was held fixed above. For the notion of topological dimension see e.g [HUREWICZ & WALLMAN].



Figure 6.1: Bifurcation within the box \mathcal{B}_{ϵ_1} for fixed small positive ϵ_1 .

6.5 Lemma: Consider system (6.1) in the open box \mathcal{B} constructed above. Let C_{per} , C_{hom} denote the period doubling and 2-hom curves, given respectively by

$$\begin{aligned} \mathcal{C}_{\text{per}} &= \{(\mathbf{r}, \epsilon) \in clos \ \mathcal{B} \mid \epsilon_2 = \kappa_{\text{per}}(\epsilon_1), \ r_0 = r_1 = \tilde{r}(\epsilon_1)\} \ , \\ \mathcal{C}_{\text{hom}}^{\epsilon_1} &= \{(\mathbf{r}, \epsilon) \in clos \ \mathcal{B}_{\epsilon_1} \mid \epsilon_2 = \kappa_{\text{hom}}(\epsilon_1), \ r_0 = 0, \ r_1 = \hat{r}(\epsilon_1) \\ or \quad r_0 = \hat{r}_1(\epsilon_1), \ r_1 = 0\} \ , \\ \mathcal{C}_{\text{hom}} &= \bigcup \ \mathcal{C}_{\text{hom}}^{\epsilon_1} \ . \end{aligned}$$

Then there exists a continuum $C \subseteq B$ of 2-per solutions of (6.1) (i.e. $r_0 \neq r_1$) with the following properties:

a)
$$clos C \cap \{(\mathbf{r}, \epsilon) \in clos \mathcal{B} \mid r_0 = r_1\} = C_{per}$$

b) $clos C \cap bdy \mathcal{B}_{\epsilon_1} = \mathcal{C}_{hom}^{\epsilon_1}, for \ 0 \le \epsilon_1 \le \epsilon_1^0$,
c) C has topological dimension
$$(6.25)$$

at least 2 at every point.

 $0 \leq \epsilon_1 \leq \epsilon_1^0$

Let C^+ denote the one point compactification of C, that is, C^+ is clos C with points on bdy $\mathcal{B} \cup C_{per} \cup C_{hom}$ identified to a single point. Then we also have

> d) there exists a continuous and essential (i.e., non-contractible) map from $clos C^+$ (6.25) to the standard 2-sphere S^2 .

Proof: The proof is basically an application of [ALEXANDER & ANTMAN, theorem 2.2]. The 1-per sheet

$$\epsilon_2 = \epsilon_2(r, \epsilon_1), r_0 = r_1 = r > 0, \epsilon_1 > 0$$

which can also be parametrized over $\epsilon = (\epsilon_1, \epsilon_2)$, plays the role of the trivial solution. Just above lemma 6.3, we have already checked the basic assumption of [ALEXANDER & ANTMAN]: a change of (Brouwer) degree viz. of sign Δ , as we cross the bifurcation curve C_{per} in the 1-per sheet. As a result, we obtain a continuum C of 2-per solutions such that (6.25.c,d) both hold. (In [ALEXANDER & ANTMAN], C is called $S_{1.}$)

To prove (6.25.a), we first claim that clos C intersects the 1-per sheet at most in the bifurcation curve C_{per} . Indeed, the linearization is nonsingular outside of C_{per} ($\Delta \neq 0$) and the implicit function theorem applies, isolating the 1-per sheet from 2-per solutions. Also, *clos* C cannot contain points in

$$\{(\mathbf{r}, \epsilon) \in bdy \mathcal{B} \mid r_0 = r_1\} \setminus \mathcal{C}_{per} .$$
(6.26)

Indeed, clos C is confined to the wedge region

$$\{(\mathbf{r},\,\epsilon) \mid \underline{\kappa}(\epsilon_1) \leq \epsilon_2 \leq \bar{\kappa}(\epsilon_1)\} \quad . \tag{6.27}$$

In this wedge, nonzero solutions with $r_0 = r_1$ and with $r_0 \in \{0, \delta^0\}$ or $\epsilon_1 \in \{0, \epsilon_1^0\}$ or $\epsilon_2 \in \{0, \epsilon_2^0\}$ do not exist. This proves our claim.

Next we claim that clos C contains C_{per} . Indeed, as we have shown above, the Brouwer degree, viz. sign Δ , changes across $\epsilon_2 = \kappa_{per}(\epsilon_1)$ for any fixed $\epsilon_1 > 0$. Therefore, $clos C \supseteq C_{per}$ by [ALEXANDER & ANTMAN, theorem 2.2]. This proves (6.25.a).

To prove (6.25.b), we first claim

$$clos \mathcal{C} \cap bdy \mathcal{B}_{\epsilon_1} \subseteq \mathcal{C}_{hom}^{\epsilon_1}$$

Indeed, the left-hand side is contained in the wedge region (6.27). But $clos C \cap bdy \mathcal{B}_{\epsilon_1}$, in that region, can consist only of 2-hom points and possibly zero. All these points are in $C_{hom}^{\epsilon_1}$.

To complete the proof of (6.25.b), we also claim

$$clos \mathcal{C} \cap bdy \mathcal{B}_{\epsilon_1} \supseteq \mathcal{C}_{hom}^{\epsilon_1}$$

Indeed, consider any ϵ_1 -section C_{ϵ_1} of C, $\epsilon_1 > 0$. By [ALEXANDER & ANTMAN, p. 349], the section C_{ϵ_1} is not contained in any compact subset of the box section \mathcal{B}_{ϵ_1} , because $clos C_{\epsilon_1}$ cannot contain any point on the 1-per sheet except for the unique bifurcation point on C_{per} with $\epsilon = (\epsilon_1, \kappa_{per}(\epsilon_1))$. Therefore,

$$clos C_{\epsilon_1} \cap bdy \mathcal{B} \neq \emptyset$$
.

In other words, $clos C_{\epsilon_1}$ contains at least one of the two points

$$\epsilon_2 = \kappa_{\text{hom}}(\epsilon_1), r_0 = 0, r_1 = \hat{r}_1(\epsilon_1)$$

or $r_0 = \hat{r}_1(\epsilon_1), r_1 = 0$

which constitute C_{hom} . By the symmetry $r_0 \leftrightarrow r_1$ of (6.1), we may then assume that $clos C_{\epsilon_1}$ also contains the other point. This proves our claim.

A minor point remains to be settled. Augmenting C_{e_1} by its symmetric counterpart, as above, could have destroyed the existence of an essential map (6.25.d), in principle. We claim this does not happen. Indeed \mathcal{C}^+ consists of three disjoint sets:

$$\mathcal{C}_0^+$$
: the collapsed point,

$$\mathcal{C}_{\boldsymbol{\zeta}}^+$$
: points in \mathcal{C} with $r_0 < r_1$,

$$\mathcal{C}^+_{\mathcal{S}}$$
: points in \mathcal{C} with $r_0 > r_1$.

Evidently, we can omit either C_{\leq}^+ or C_{\geq}^+ from C^+ and still find an essential map into S^2 ; say

 $\mathcal{C}^+_0 \ \cup \ \mathcal{C}^+_< \longrightarrow S^2$.

Extending this map to $\mathcal{C}^+_>$ such that orbits under $r_0 \longleftrightarrow r_1$ get mapped to the same point, we find a symmetric continuum \mathcal{C} for which (6.25.d) still holds.

This completes the proof of lemma 6.5, and the proof of theorem B.

7. Discussion

We put our main theorems A, B in a variety of perspectives. First we compare and contrast our results, as they stand, with earlier work by other authors. Then we re-emphasize a central weakness of our approach via Ljapunov-Schmidt reduction: we have lost control over N-hom/N-per orbits with N > 2. We indicate, how this defect can be remedied by constructing a suitable center manifold. Because our discussion of Poincaré return maps has caused us a lot of technical troubles, so far, we also explain why another idea, which is based on exponential dichotomies, does not work as well for homoclinic bifurcations. Hard core applications of our results are very scarce, so far. We relate this deplorable fact to the lack of numerical pathfollowing schemes for homoclinic orbits. After pointing out analytical difficulties with our results in infinite-dimensional settings, we return to the pathfollowing question from a global, theoretical point of view. We sketch an idea for an emerging index theory for paths of homoclinic orbits which hopefully should bridge the gap between local "birth" of homoclinic orbits at B-points and, on the other side, chaotic dynamics of the Shilnikov type. We summarize some known ingredients, but much more still remains to be done.

Postponing further speculation, for a little while, let us return to the homoclinic side-switching bifurcation. According to theorem A, this bifurcation is associated to non-twisted homoclinic orbits and can therefore occur for vector fields in the plane. Indeed, [SANDERS & CUSHMAN] (1984) have found resonant side-switching to occur in the (autonomous) equation for a Josephson junction

$$\dot{\phi} = y$$

$$\dot{y} = -\sin\phi + \epsilon(a - (1 + \gamma\cos\phi)y) ,$$
(7.1)

with $\phi \in S^1$, ϵ small positive, and with real parameters a, γ . They obtain resonant side-switching near $a = 16/3\pi$, $\gamma = 1$. In particular, the correct asymptotic expression for the per-fold, given in (1.16), was obtained in [SANDERS & CUSHMAN, (5.11)]. This includes the computation of a_0 and of the finite limit in (1.16). Their explicit computations are based on averaging of a perturbed pendulum equation, Picard-Fuchs equations, and associated Riccati equations. On the other hand, our much less explicit approach carries over to vector fields which are far from a Hamiltonian structure and to vector fields in higher space dimensions. An earlier reference which is closely related to resonant side-switching is [LEON-TOVICH] (1951). There, it is stated in theorem 3 (putting n = 0, $c_0 = 0$, $a_1 \neq 0$) that up to two periodic orbits can co-exist for perturbations of a resonant homoclinic orbit in the plane. Still for planar C^2 vector fields, [NOZDRACHEVA] (1979) essentially obtained theorem A. Tangency of the per-fold at $\epsilon_1 = 0$ was noticed, but the asymptotic expression (1.16) was not derived.

The bifurcation of 2-hom orbits is a question which was first addressed by [YANAGIDA]

(1986) in a travelling wave context as follows. Let

$$\zeta_t = D\zeta_{xx} + G(\alpha_2, \zeta), \ x \in \mathbb{R}, \ \alpha_2 \in \mathbb{R}$$
(7.2)

be a reaction diffusion system with nonnegative diagonal diffusion matrix D and with nonlinearity G. Then special solutions of the form $\zeta = \zeta(x + ct)$, called travelling waves, satisfy the ODE

$$0 = -c\dot{\zeta} + D\ddot{\zeta} + G(\alpha_2, \zeta) , \qquad (7.3)$$

which is of the form

$$\dot{z} = F(\alpha, \zeta) , \qquad (1.1_{\alpha})$$

with suitably defined z and with $\alpha = (\alpha_1, \alpha_1) := (c, \alpha_2)$. Specifically, Yanagida considers 2-hom orbits to an equilibrium with unstable dimension 1 (n = 1, in ournotation). This assumption enables him to treat the Poincaré map directly, avoiding Shilnikov variables like (s, x^{in}, y^{out}) . Unfortunately, he completely linearizes F near the equilibrium z = 0 and then omits terms which are of higher order, formally. Linearizations of class C^k , $(k \ge 1$ to preserve tangencies), require certain diophantine nonresonance conditions on the eigenvalues. For linearization of flows which do not contain parameter, see e.g. [BELITSKII, STERNBERG 1-2, SELL 1-2. In particular, our system violates the Sternberg condition of order 3 as soon as $z \in \mathbb{R}$, due to resonance. For smooth normal forms in the plane at resonance see [BOGDANOV 3]. We point out that the linearization question is related to the exponential expansion (3.9), at least for $z \in \mathbb{R}^2$; see [DENG1, section 6]. Anyway, Yanagida derives the correct (truncated) reduced normal form equation for 2-hom orbits, concluding existence in the twisted and nonexistence in the non-twisted case. We have now confirmed these results, extending them to higher unstable dimension and deriving exponential asymptotics of the bifurcation curve κ_{hom} . As we have mentioned in section 2 above, [GLENDINNING 2] (1987) has computed both the asymptotics for $\kappa_{\rm hom}$ and $\kappa_{\rm per}$ for the truncated reduced normal form. He claims that reduction to this truncated form can be achieved "via standard techniques". In a way, we have shown that he is right: a posteriori, and modulo some subtleties described below. We note here already that even the discussion of the reduced normal form equation (4.13) is a nontrivial task, since a systematic theory of equations where parameters enter in the exponents, i.e. transcendentally, does not exist. In particular, it is not clear how to neglect terms "of higher order". [KOKUBU 1] (1987) also notes the linearization flaw in Yanagida's presentation, announcing an alternative proof. Accidentally, the crucial twist condition is missing in his statement of Yanagida's result [KOKUBU 1, theorem 3]. Writing this discussion, we obtain the preprint [KOKUBU 2] (1988) which, in theorem D, gives a correct version of Yanagida's result in three space dimensions (n = 1, m = 2). The proof avoids linearization and relies on the Shilnikov expansion given in proposition 3.2 above. The method of proof differs from ours. In particular, periodic orbits are not discussed and the homoclinic orbits are not obtained through a Ljapunov-Schmidt reduction. Instead, a transversality assumption is made. Specifically, the existence of an (m + 1)-dimensional invariant manifold W^{su} is used, which is assumed to be tangent to $T_0W^s \oplus \text{span}(e^+)$ at z = 0. A manifold W^{us} , tangent to $T_0W^u \oplus \text{span}(e^-)$, can be defined analogously. Transversality is assumed for the intersections

$$W^u \cap W^{su}, W^s \cap W^{us}$$

This assumption enters already into continuation of the primary 1-hom branch. Existence of W^{su} , W^{us} entails a loss of regularity. These manifolds can be assumed to exist of class C^k , if the spectrum of the linearization at z = 0 satisfies the gap condition

$$\min\left(\bar{\mu}_{1}, \bar{\nu}_{1}\right) > k \cdot \max\left(\mu_{0}, \nu_{0}\right)$$

in the notation of assumption (1.6); see [HIRSCH & PUGH & SHUB, section 5, VANDERBAUWHEDE]. Generically, this forces us down to C^1 -regularity throughout. In contrast, we obtain a 2-hom curve κ_{hom} which is smooth except at bifurcation. [KOKUBU 2] notices tangency of κ_{hom} to the 1-hom curve but does not derive the exponential expansion

$$0 < \lim_{\epsilon_1 \searrow 0} \kappa_{\text{hom}}(\epsilon_1) |a_0|^{1/\epsilon_1} < \infty$$
.

However, the main thrust in [KOKUBU 2] is on heteroclinic phenomena. This explains why periodic orbits are not considered.

We emphasize that Yanagida has, at least formally, found two other generic mechanisms for non-resonant homoclinic doubling besides the resonant case, assuming $\mu_0 > \nu_0$; see [YANAGIDA, theorem 3.3]. Geometrically, these other two mechanisms work as follows. Suppose we follow a path of homoclinic orbits $z^*(t)$, varying the parameter α along a curve. Then,

$$\lim_{t \to +\infty} \dot{z}^*(t) / |\dot{z}^*(t)|$$

might switch from the principal stable eigenvector $-e^-$ to e^- itself, somewhere along this curve. In between, there will be a parameter value, viz. a bifurcation point, where the above limit is a non-principal eigenvector. This is the first mechanism. Alternatively, the unstable manifold W^u can intersect W_{loc}^s at the point q in an exceptional position, so that the strong λ -Lemma 3.3 does not apply. In other words, the status of the homoclinic orbit $z^*(t)$ switches from twisted to nontwisted at some (bifurcation) point along the parameter curve. This is the second mechanism which leads to homoclinic doubling. Both mechanisms are discussed under the, hopefully unnecessary, assumption n = 1 for the unstable dimension. At this stage, it seems likely that all generic mechanisms for homoclinic doubling involving real principal eigenvalues have been found. Strictly speaking, however, we are still lacking rigorous necessary criteria for homoclinic bifurcations. For reasons of space, we cannot pursue these questions any further here. It is a central weakness of our approach that we lose control over N-per/N-hom orbits, $N \ge 3$, in our analysis of the reduced normal form

$$r_{j+1} = \epsilon_2 + a(\epsilon)r_j^{1+\epsilon_1} + O(|\mathbf{r}|^{1+\omega}), \ j \pmod{N} , \qquad (4.13)$$

for $\mathbf{r} = (r_0, \ldots, r_{N-1})$. As we have noticed in section 2, the *O*-term is the reason why (4.13) does not describe an iteration of a single monotone scalar function, for which *N*-per/*N*-hom orbits with $N \ge 3$ could not occur. This subtlety seems to have escaped the attention of some authors. We briefly describe two different angles of attack to the problem.

First, we can approach the problem algebraically. For example, consider the nontwisted case a > 0, viz. a > 1. For N = 2, it can be shown that no additional solutions occur besides those with $r_0 = r_1 = r$, which are seen for N = 1 already. Indeed, suppose $0 \le r_0 < r_1$. Then we can reconsider the proof of lemma 6.4, this time for a > 1 rather than a < -1. We obtain a contradiction from (6.22), since both a and σ_{ϵ_1} are positive. For $N \ge 3$, however, these essentially scalar convexity arguments fail. In particular, we are not able to gain sufficient control over (4.13) to exclude the existence of N-per solutions algebraically.

Second, let us approach the problem more geometrically. Suppose there exists a two-dimensional center manifold attached to the homoclinic orbit $z^*(t)$, which describes bifurcations just as in the case of a periodic orbit. In a Poincaré section, we then have a one-dimensional invertible return map. Thus N-per orbits with $N \ge 3$ cannot occur nearby, since they would have to lie on the center manifold. But: does there actually exist such a center manifold W^c ? We sketch how W^c might be constructed, at least of class C^1 , following an idea which was pointed out to us by J. Guckenheimer. Basically, one would like to use the (m+1)-dimensional manifold W^{su} , tangent to $T_0W^s \oplus \text{span}(e^-)$, which we have mentioned above in the context of [KOKUBU 2]. Just assume that W^{su} and W^{us} intersect transversely along the orbit $\Gamma = \{z^*(t)\}$ at $\alpha = 0$ and define

$$W^c = W^{su} \cap W^{us} ;$$

obviously dim $W^c = 2$. If W^{su} and W^{us} are constructed very carefully, then W^c will contain all orbits which remain in a tubular neighborhood U of the homoclinic orbit Γ . In particular, all N-hom and N-per orbits will lie in W^c , as required above. For a similar approach to a simplified model of the Lorenz system see [ROBINSON].

Constructing W^{su} , W^{us} for this purpose involves the following subtlety. The vector field F has to be modified such that W^{su} , e.g., can be constructed globally to consist of all those trajectories which do not escape to infinity at a rate faster than

$$e^{(\nu_0+\epsilon)t}$$
, $t\to\infty$.

Usually, F is allowed to be modified everywhere outside a small neighborhood of the origin [HIRSCH & PUGH & SHUB, section 5A, VANDERBAUWHEDE]. For our

purposes, however, F must not be modified in a tubular neighborhood U of the entire homoclinic loop Γ . Therefore, constructing W^c will be a not so standard task, involving a global extension of the vector field to an appropriately chosen bundle over the homoclinic loop Γ . Once W^c is shown to exist, there remains a question about (linearized) stability of the bifurcating 1-per and 2-per solutions. We hope that a "principle of reduced stability" in the sense of [KIELHÖFER & LAUTERBACH] holds: this would allow us to determine stability of the periodic solutions directly from the reduced normal form (4.13).

An alternative approach to pathfollowing of homoclinic orbits, based on exponential dichotomies, can be sketched as follows. We rewrite (1.1) as

$$\mathcal{F}(\alpha, z(\cdot)) := -\dot{z}(t) + F(\alpha, z(t)) = 0 ,$$

$$\mathcal{F} : \mathbb{R}^2 \times Z_1 \longrightarrow Z_0$$
(7.4)

where Z_{ι} , $\iota = 0, 1$, are suitably chosen Banach spaces like $BC^{\iota}(\mathbb{R}, \mathbb{R}^{m+n})$ with the uniform sup-norm, or like the Sobolev spaces $H^{\iota}(\mathbb{R}, \mathbb{R}^{m+n})$. It follows from [PALMER, lemma 4.2] that the linearization

$$D_z \mathcal{F}(0, z^*(\cdot))$$

at the homoclinic orbit $z^*(\cdot)$ is a Fredholm operator of Fredholm index zero. The results in [PALMER] are based on exponential dichotomies of certain families of projection operators along the orbit $z^*(t)$. Note that $\dot{z}^*(\cdot)$ is always in the kernel of $D_z \mathcal{F}(0, z^*(\cdot))$. Assuming surjectivity of the total derivative $D \mathcal{F}$ makes pathfollowing of homoclinic orbits amenable to an application of the implicit function theorem, and even to global continuation techniques. The formulation (7.4) has a fundamental drawback, however, when it comes to bifurcation problems. For illustration, consider resonant homoclinic doubling (theorem B). As $\epsilon_1 \searrow 0$, $\epsilon_2 = k_{\text{hom}}(\epsilon_1)$, the 2-hom trajectory $\{\hat{z}(t) \mid t \in \mathbb{R}\}$ tends to the 1-hom trajectory $\{z^*(t) \mid t \in \mathbb{R}\}$ in phase space $z \in \mathbb{R}^{m+n}$. In contrast, $\hat{z}(\cdot)$ does not tend to $z^*(\cdot)$ in any of the function spaces Z_0 or Z_1 . In fact, $\hat{z}(\cdot)$ does not have a limit for $\epsilon_1 \searrow 0$. A similar obstacle arises, when we try to capture periodic orbits limiting on a 1-hom at a blue sky catastrophe in the functional analytic framework of (7.4).

Hard core applications of our results are scarce. Resonant side-switching, as we have mentioned above, was found by [SANDERS & CUSHMAN] (1984) to occur in the Josephson junction. [YANAGIDA] (1986) presents his result in the framework of a "generalized nerve equation introduced by [FITZHUGH]", without being entirely specific about the underlying nonlinearities. Actually, this situation is somewhat reminiscent of period doubling, which is hardly ever detected analytically for a concrete given system. Rather, there are numerical pathfollowing routines for periodic orbits which find period doublings in a broad variety of

applications. See for example [DOEDEL & KERNEVEZ, MAREK & KUBIČEK, SEYDEL]. Efficient pathfollowing codes for homoclinic orbits in two parameters do not exist. In AUTO [DOEDEL & KERNEVEZ], homoclinic orbits are treated as periodic orbits of "large" period. The special structure of homoclinic orbits near stationary solutions is thus ignored. Concerning the numerical treatment of one-sided asymptotic boundary value problems we mention [de HOOG & WEISS, LENTINI & KELLER]. Note that, in the functional analytic framework of (7.4) above, a homoclinic orbit $z^*(t)$ solves the boundary value problem (7.4) with the two-sided asymptotic boundary condition $z^*(t) \rightarrow 0$ for $t \rightarrow \pm \infty$. Of course, this setting suffers from the above mentioned difficulties at bifurcations. The exponential asymptotics (1.16), (1.18-19), (5.9), (6.7), (6.15) pose an even more aggravating obstacle. For example, consider resonant homoclinic doubling (theorem B). Then the 2-hom branch is close of order

$$|a_0|^{-1/\epsilon_1} , \ a_0 < -1 , \tag{7.5}$$

both in parameter and in phase space, as we move away from the bifurcation point along the 1-hom branch by an arclength of $\epsilon_1 > 0$. For moderate $a_0 = -2$, $\epsilon_1 = 0.04$, the quantity (7.5) is still as small as $3 \cdot 10^{-8}$. Such effects are hard to detect in numerical case studies. For completeness, we mention that 2-hom orbits were detected analytically in a FitzHugh-Nagumo system with a complex pair of principal eigenvalues rather than real eigenvalues; see [EVANS & FENICHEL & FEROE, FEROE, HASTINGS] (1982). This is related to a celebrated result by [SHILNIKOV 2] (1964), predicting shift dynamics near certain homoclinic orbits of flows. We return to this phenomenon at the end of this section.

One potential class of applications is concerned with infinite-dimensional dynamical systems. Specifically, we think of reaction diffusion systems, viz. analytic semigroups, and of differential equations involving time delays. We caution the reader, that it is not clear at present how to carry over the fundamental expansion

$$y^{\text{in}} = r(\psi + R_y)$$

$$x^{\text{out}} = r^{1/\nu_0}(\varphi + R_x)$$
(3.9)

to such infinite-dimensional semiflows. In fact, the proof of (3.9) involves flows in both positive and negative time direction; see [DENG 1]. Therefore, we do not claim that our results hold, e.g., for reaction-diffusion equations. Techniques are being developed, though, to study homoclinic and heteroclinic orbits for infinitedimensional semiflows in a more systematic fashion. We mention results on the blue sky catastrophe and an infinite-dimensional λ -Lemma [CHOW & DENG], and the results by [LIN, HALE & LIN] involving an exponential dichotomy setting as in (7.4) for functional differential equations. For inclination lemmas and bifurcations of homoclinic/heteroclinic orbits for functional differential equations see also the fundamental papers [WALTHER 1-5]. For a detailed and explicit investigation of heteroclinic orbits of scalar reaction diffusion equations in one space dimension see [BRUNOVSKÝ & FIEDLER 2-3, ANGENENT & FIEDLER] and the references there. A scalar delay equation was treated in [FIEDLER & MALLET-PARET], in a similar spirit. Representing an entirely different line of thought, we finally mention [KIRCHGÄSSNER 1-3, FISCHER] where homoclinic orbits of small amplitude are detected via a reduction to a finite-dimensional center manifold. These results essentially deal with PDEs of elliptic type in unbounded domains like strips or cylinders. The homoclinic structure arises via appropriate boundary conditions at infinity in a setting which is an infinite dimensional version of (7.4), even though an elliptic system does not define a dynamical system, a priori. Time in (7.4) gets replaced by the unbounded space coordinate of the elliptic system. For the "nonautonomous" case, in this sense, see [KIRCHGÄSSNER 2-3, MIELKE 1-2] and the references there. Global continuation of homoclinic orbits is possible in principle, see e.g [KIRCHGÄSSNER, AMICK & KIRCHGÄSSNER], but suffers from the inherent drawbacks of the setting (7.4) which we have described above.

Returning to finite-dimensional questions, we mention the problem of nonautonomous perturbations. For example, consider

$$\dot{z}(t) = F(\alpha, z(t)) + \epsilon h(t)$$
 $\alpha = (\alpha_1, \alpha_2)$,

for periodic forcing h with small amplitude $|\epsilon|$. Fixing $\epsilon \neq 0$ small, it is a difficult , question to study the dynamic behavior of the corresponding two-parameter family of period map diffeomorphisms, as α varies in a neighborhood of a (non-twisted or twisted) homoclinic bifurcation associated to $\epsilon = 0$. For an example concerning a homoclinic orbit hitting a stationary fold in the Josephson equation as in figure 7.3 below, see [SCHECTER 2].

Hamiltonian systems and time reversible systems are other classes of equations which we have excluded here. It is a common feature of these systems that any linearization at any equilibrium has symmetric spectrum, i.e.

$$\mu \in \operatorname{spec} \iff -\mu \in \operatorname{spec}$$
.

In particular, any homoclinic orbit with real principal eigenvalues is automatically resonant in our sense (1.9). In this context, we remember that resonant sideswitching was found by [SANDERS & CUSHMAN], perturbing the (Hamiltonian) mathematical pendulum. We are not aware of any results on resonant homoclinic doubling for Hamiltonian or reversible systems. For a discussion of certain Hamiltonian systems with periodic orbits, homoclinic orbits, and heteroclinic loops see [HOFER & TOLAND].

After these excursions, let us now reconsider our original generic two-parameter problem

$$\dot{z} = F(\alpha, z) , \ \alpha \in \mathbb{R}^2 .$$
 (1.1_{\alpha})

Broadening our view, in the remainder of this section, we address some questions about global pathfollowing of homoclinic orbits. As a starting point, we resume our discussion of *B*-points, alias the Arnold-Bogdanov-Takens singularity, from section 1 (see (1.4) and figure 1.1). At a *B*-point, as we recall, a path of homoclinic orbits starts (or terminates). As a goal, we would like to define an orientation of these paths in parameter space, globally, such that we can follow oriented paths "as far as possible" without ever cycling back onto a previously followed piece of the path. Instead of rendering this goal artificially precise, we now define an orientation. Then we will see, what this does for us. Our attempt is much in the spirit of [MALLET-PARET & YORKE 1,2], [ALLIGOOD & MALLET-PARET & YORKE], [CHOW & MALLET-PARET & YORKE], [FIEDLER 1,2].

As a prerequisite, we define an index $\Phi \in \{-1, 0, +1\}$ for a hyperbolic periodic orbit $\gamma = \{z(t) \mid 0 \leq t \leq p\}$ of minimal period p > 0. Hyperbolic means that γ does not possess Floquet multipliers on the unit circle, except the trivial one. Let σ^+, σ^- denote the number of real Floquet multipliers in the intervals $(1, \infty), (-\infty, -1)$, respectively, counting algebraic multiplicities. Then the orbit index Φ is defined as

$$\Phi := \frac{1}{2}((-1)^{\sigma^+} + (-1)^{\sigma^+ + \sigma^-}) \quad . \tag{7.6}$$

This orbit index Φ , defined first in [MALLET-PARET & YORKE 1] (1979), turns out to be a local homotopy invariant for periodic orbits of generic one-parameter vector fields.

Equipped with the orbit index Φ , we can now define an orientation of certain homoclinic paths in parameter space. We assume that periodic orbits γ near the homoclinic orbits occur only for parameters α on one side of the local path segment which we consider (see figure 7.1). We then orient the homoclinic path segment such that the periodic orbits γ are on the right if $\Phi = +1$. If $\Phi = -1$, then we want γ to be on the left. If $\Phi = 0$, then we refuse to define an orientation.



Figure 7.1: Orienting homoclinic paths

How compatible is this definition with the local bifurcation picture at a *B*-point? Note that $\sigma^- = 0$ and hence $\Phi \neq 0$, near a *B*-point. Therefore, the homoclinic path can be oriented. Following its orientation, the homoclinic path can either emanate from the *B*-point, or it can terminate there. These two cases fit nicely with the index *B* of *B*-points, which was defined in [FIEDLER 2] for the purpose of global Hopf bifurcation. Specifically, B = +1 when a homoclinic path emanates and B = -1 at termination. The index *B* is defined in local terms and is easily accessible numerically. See the code BALCON [FIEDLER & KUNKEL 1,2] which is based on ALCON [DEUFLHARD & FIEDLER & KUNKEL].

What happens to the orientation, far from *B*-points? Consider resonant sideswitching, first. By theorem A, figure 1.3.a, there are two 1-per solutions at parameters α in the region II. Their respective orbit indices Φ are opposite in sign (or are both zero), by homotopy invariance near the fold curve $\epsilon_2 = \kappa(\epsilon_1)$. One of the periodic orbits continues to a blue sky catastrophe at the right 1-hom branch $\epsilon_1 > 0$, $\epsilon_2 = 0$. The other orbit continues to the left 1-hom branch $\epsilon_1 < 0$, $\epsilon_2 = 0$. Since the 1-per orbits have opposite index Φ and approach the 1-hom branch from opposite sides, the orientation of the 1-hom branch is defined consistently on both sides of the bifurcation point $\epsilon = 0$.

As a second example, consider resonant homoclinic doubling (theorem B and figure 1.3.b). To determine all orientations, fix ϵ_2 small positive and vary only ϵ_1 . Crossing the κ_{per} -region in the 1-per sheet from left to right, we see that the number σ^- of Floquet-multipliers in $(-\infty, -1)$ changes by one. Thus $\Phi = 0$ on one side,

and $\Phi \neq 0$ on the other side. Suppose $\Phi = +1$, for definiteness, as in figure 7.2. For simplicity, also suppose that the 2-per sheet is confined to the region IV and is parametrized over ϵ . Then Φ on the 2-per sheet can be computed, by homotopy invariance of Φ , near the period doubling curve $\epsilon_2 = \kappa_{per}(\epsilon_1)$.



Figure 7.2: Two cases of resonant side-switching

If no further period doublings occur on the 2-per sheet (leading to 4-per solutions) then Φ extends, constantly, to the 2-hom branch. This defines an orientation of the 2-hom branch, as in figure 7.2. We conclude that, in each case, precisely two of the three homoclinic half-branches are oriented, and this orientation extends consistently through the bifurcation point. These arguments can be extended to cover the case when the 2-per sheet is neither confined to the region IV nor parametrized over ϵ . However, due to the difficulties which we have described above, we cannot exclude the possibility of further period doublings from the 2-per sheet which would lead to 4-per orbits. In fact, we are lacking control over N-per orbits with $N \geq 3$. This difficulty does not arise for resonant side-switching because 2-per orbits do not exist, in that case.

Of course, the above remarks do not constitute a global bifurcation result for paths

of homoclinic orbits. Other bifurcations can occur. We briefly mention some of the known results. But we do not tie these results in with our index theory, this time.

We have already mentioned above that non-resonant mechanisms for homoclinic doubling were found by [YANAGIDA]. Now let us consider the base point A of a homoclinic orbit $z^*(t)$, that is, the stationary point to which $z^*(t)$ tends for $t \rightarrow \pm \infty$. In the previous sections we have discussed at length what happens when the principal eigenvalues at A become resonant. Now suppose one of the principal eigenvalues becomes zero. Generically, this means that the homoclinic path hits a fold line in parameter space while the base point A hits the fold. This situation was studied by [LUKYANOV] (1979) and [SCHECTER 1] (1985) for $z \in \mathbb{R}^2$. The bifurcation diagram is sketched in figure 7.3. It was noted by [SCHECTER] that the hom-curve hits the fold curve tangentially rather than transversely. Corresponding results for higher dimension of z are now avialable; see [CHOW & LIN, DENG] (1988). Also in higher space dimension, but in one-parameter systems, the generation of (unique) periodic orbits from homoclinic orbits with base point on a fold was studied by [SHILNIKOV 1,3] (1962, 1966). [SHILNIKOV 6, AFRAIMOVICH & SHILNIKOV (1969, 1974) also have observed that a fold point A, with $p \geq 2$ distinct homoclinic orbits attached to it, generates nearby dynamics which is equivalent to shift dynamics on p symbols. Evidently, this situation can arise during a pathfollowing process when p path segments successively hit a stationary fold, each time as in figure 7.3, and then co-exist. In other words, the very intriguing results by [SHILNIKOV 6, AFRAIMOVICH & SHILNIKOV] indicate how pathfollowing of homoclinic orbits can lead to regions with chaotic, i.e. shift-type, dynamics near a stationary fold.



Figure 7.3: A homoclinic path hitting a stationary fold.

At this stage, it seems natural to ask what happens when the base point approaches a Hopf bifurcation along the homoclinic path. Long before that can happen, however, another celebrated result by Shilnikov takes effect. Namely, we may assume that both principal eigenvalues are in fact conjugate complex pairs, or that one of them is real, say $\nu_0 > 0$, while the other "one" is a conjugate complex pair, say $-\mu_0 \pm i\omega_0$, such that

$$0 < \mu_0 < \nu_0 \quad . \tag{7.7}$$

In both cases, nearby shift dynamics on a countable alphabet was discovered, generically. See [SHILNIKOV 2,4,7] (1965, 1967, 1970). A discussion of some of these results is given in [DENG 3] (1988), with an emphasis on exponential expansions of the Shilnikov variables like (3.9). For a C^1 -version of (7.7) with $z \in \mathbb{R}^3$ see also [TRESSER] (1984). For applications of these ideas in the FitzHugh-Nagumo equations see [EVANS & FENICHEL & FEROE, FEROE, HASTINGS] (1982), again. Homoclinic tangencies of return maps occur on a dense set along homoclinic paths where (7.7) holds, see [OVSYANNIKOV & SHILNIKOV] (1986). The transition into a region where (7.7) holds was studied for two parameter vector fields in \mathbb{R}^3 by [BELYAKOV 1,2] (1980, 1984). He detects, e.g., countably many paths of 2-hom and, in one of the cases, of 3-hom orbits accumulating at the transition point. See also [RODRIGUEZ] (1986), for the 1- hom branch.

When (7.7) is violated, that is

$$0 < \nu_0 < \mu_0$$
,

then a 2-shift still occurs in \mathbb{R}^3 , provided that two distinct homoclinic orbits are limiting onto the same base point A, thus forming a figure "8". The two symbols correspond to excursions along the upper/lower loop of the figure "8". In particular, such a situation can arise for systems with a (reflection) \mathbb{Z}_2 -symmetry; see [HOLMES, TRESSER, GAMBAUDO & GLENDINNING & TRESSER] (1980, 1984). The case $\nu_0 > \mu > 0$ was also discussed by [TRESSER, GLENDINNING 1] (1984). Repeated period doublings near pairs of homoclinic orbits with real principal eigenvalues and coinciding principal directions

$$\lim_{t\to+\omega} \dot{z}^*(t)/|\dot{z}^*(t)| ,$$

due to \mathbb{Z}_2 -symmetry, are observed by [ARNEODO & COULLET & TRESSER, KU-RAMOTO & KOGA, COULLET & GAMBAUDO & TRESSER, LYUBIMOV & ZAKS] (1980, 1982–1984). For the Lorenz system, see in particular [SPARROW]. Again, complicated dynamics abound.

Towards the end of our long excursion into global pathfollowing, let us mention the largely open problem of bifurcation from heteroclinic cycles. From a pathfollowing point of view, heteroclinic cycles arise as follows. Let A denote the base point of a homoclinic orbit Γ , as before. Following a path in parameter space, some part of Γ might approach another stationary point **B**. In the limit, we have a heteroclinic cycle: one trajectory runs from A to B and another one returns from B to A. As references for bifurcations from heteroclinic cycles one may consult [REYN] for $z \in \mathbb{R}^2$, [BYKOV, TRESSER] (1980, 1984) for $z \in \mathbb{R}^3$, when conjugate complex eigenvalues occur, and [RINZEL & TERMAN, CHOW & DENG & TERMAN 1-2, DENG 2, KOKUBU 1-2] (1981, 1986-1988) when the principal eigenvalues are real. In all these cases, it is assumed that both A and B are hyperbolic with unstable dimension given by $u_A = u_B = 1$. In some cases, [BYKOV, TRESSER] find shift-dynamics of the Shilnikov type. [DENG 2], on the other hand, detects a phenomenon where, in parameter space, paths $\kappa_0, \kappa_1, \ldots$ of heteroclinic solutions bifurcate from the heteroclinic cycle point such that the following holds. For $\alpha \in \kappa_N$, the heteroclinic orbit cycles through N complete loops near the heteroclinic cycle and only afterwards converges to A, B for $t \to \pm \infty$ without cycling further. The basic assumption there is again a certain twist condition. [DENG 2] also sketches how this bifurcation can occur in planar vector fields. In general, heteroclinic cycles break apart under perturbations of the vector field; generically they occur as a phenomenon of codimension at least two. When enough symmetry is present, however, heteroclinic cycles can become structurally stable. Essentially, heteroclinic orbits can then come to lie in certain linear subspaces which have a symmetry-invariance and are therefore flow invariant. Of course, this influences the local bifurcation diagrams as well. For interesting examples see [ARMBRUSTER & GUCKENHEIMER & HOLMES, GUCKENHEIMER & HOLMES] (1988). But even when symmetry is not present, there remains an enormous number of largely unresolved cases of generic two-parameter bifurcations from heteroclinic cycles. These cases arise when the unstable dimensions u_A or u_B exceed one, possibly differing from each other, with real or complex principal stable or unstable eigenvalues of A and **B**. For an intricate example of heteroclinic loops in the \mathbb{Z}_2 -symmetric Lorenz system see [GLENDINNING & SPARROW] and also [SPARROW].

Each of the above cases, separately, will create a challenge to our attempt of sketching an emerging theory for global pathfollowing of homoclinic orbits in twoparameter flows. On this long way still ahead, it is our firm hope that paths of homoclinic orbits will turn into a systematic guiding thread from easily detected B-points up to regions with complicated, shift-type dynamics.

8. Appendix: genericity assumptions

In this appendix, we summarize the nondegeneracy assumptions which enter into our main theorems A and B.

Our first assumption was

$$\operatorname{codim} (T_{\mathbf{p}}W^{u} + T_{\mathbf{p}}W^{s}) = 1 , \qquad (1.10.a)$$

so that $T_{\mathbf{p}}W^s \cap T_{\mathbf{p}}W^u$ is given by the span of the tangent $\dot{z}^*(0)$ of the homoclinic orbit at $z^*(0) = \mathbf{p}$, $\alpha = 0$. Then we have assumed

$$\mathbf{p} \notin W^{ss}$$
, $\mathbf{p} \notin W^{uu}$, (1.10.b)

so that $z^*(t)$ approaches z = 0 tangentially to the principal eigendirections e^{\mp} as $t \to \pm \infty$. With the notation

$$T_{z^*(t)} := T_{z^*(t)} W^s + T_{z^*(t)} W^u ,$$

we have then assumed that $T_{z^*(t_0)}$, say, avoids certain exceptional positions. In that case, the strong λ -Lemma (proposition 3.3) implies

$$\lim_{t \to -\infty} T_{z^{\star}(t)} = T_0 W^{ss} \oplus T_0 W^u ,$$

$$\lim_{t \to +\infty} T_{z^{\star}(t)} = T_0 W^s \oplus T_0 W^{uu} .$$
(1.13)

This assumption has enabled us to discern twisted and non-twisted homoclinic orbits. Moreover, this assumption made our Ljapunov-Schmidt reduction work, via the projection P_i^0 defined in (4.6).

So far, our genericity assumptions are concerned only with the vector field $F = F(\alpha, z)$ for fixed parameter $\alpha = 0$. These assumptions led to the reduced bifurcation equation

$$\Phi_{j}(\mathbf{r}, \alpha) = -c_{0}(\alpha) + c_{1}(\alpha)r_{j+1} - c_{2}(\alpha)r_{j}^{1/\nu_{0}(\alpha)} + O(|\mathbf{r}|^{1+\omega}) , \ j(\text{mod } N) ,$$
(4.8)

with appropriate coefficients $c_0(\alpha)$, $c_1(\alpha)$, $c_2(\alpha)$, given explicitly by (4.9). The remaining genericity assumptions enable us to rewrite (4.8) as the reduced normal form

$$r_{j+1} = \epsilon_2 + a(\epsilon) r_j^{1+\epsilon_1} + O(|\mathbf{r}|^{1+\omega}) , \ j \pmod{N} , \qquad (4.13)$$

with $|a(0)| \neq 0, 1$, by a parameter transformation

$$\alpha \to \epsilon = \epsilon(\alpha)$$
.

This diffeomorphism is given explicitly in (4.14.a,b). Denoting the principal unstable eigenvalue by $\nu_0 = \nu_0(\alpha)$, with the principal stable eigenvalue normalized to be $-\mu_0(\alpha) = -1$, the relevant nondegeneracy assumptions were:

a) $D_{lpha} c_0(0)$ and $D_{lpha} u_0(0)$ are

linearly independent,

b)
$$c_1(0) > 0$$
, (4.15)
c) $\left| \frac{c_2(\alpha)}{c_1(\alpha)} \right| \neq 0, 1$.

Genericity of these three assumptions was discussed in the proof of corollary 4.3. In particular positivity of $c_1(0)$ was generic, due to a sign convention.

With (1.10.a-b), (1.13) and (4.15.a-c), our list of genericity assumptions is complete.

Acknowledgement

One of us, B. Fiedler, has been supported by a Konrad Zuse Fellowship. Moreover, he is indebted to J. Alexander, R. Bogdanov, P. Deuflhard, J. Guckenheimer, G. Iooss, and H.-O. Walther for helpful discussions. Special thanks go to E. C. Körnig for typesetting the whole manuscript in $T_{\rm E}X$ with extreme care and insight.

.
References:

- **R. Abraham & J. Marsden:** Foundation of Mechanics. Benjamin/Cummings, Reading (1978).
- R. Abraham & J. Robbin: Transversal Mappings and Flows. Benjamin Inc., Amsterdam (1967).
- V.S. Afraimovich & L.P. Shilnikov: On attainable transitions from Morse-Smale systems to systems with many periodic points. Math. USSR Izvestija 8, pp. 1235 - 1270 (1974).
- J.C. Alexander & S.S. Antman: Global and local behavior of bifurcating multidimensional continua of solutions for multiparameter nonlinear eigenvalue problems. Arch. Rational Mech. Analysis 76, pp. 339-354 (1981).
- K.T. Alligood & J. Mallet-Paret & J.A. Yorke: An index for the global continuation of relatively isolated sets of periodic orbits. In [PALIS], pp. 1 – 21.
- C.J. Amick & K. Kirchgässner: A theory of solitary water-waves in the presence of surface tension. Preprint (1988).
- S. Angenent & B. Fiedler: The dynamics of rotating waves in scalar reaction diffusion equations. Trans. AMS 307, pp. 545 568 (1988).
- D. Armbruster & J. Guckenheimer & P. Holmes: Heteroclinic cycles and modulated travelling waves in systems with O(2) symmetry. Physica 29D, pp. 257 - 282 (1988).
- A. Arneodo & P. Coullet & C. Tresser: A possible new mechanism for the onset of turbulence. Phys. Lett. 81 A, pp. 197 201 (1981).
- V.I. Arnol'd: Lectures on bifurcations and versal systems. Russ. Math. Surv. 27, pp. 54 123 (1972).
- G.R. Belitskii: Functional equations and conjugacy of local diffeomorphisms of a finite smoothness class. Funct. Analysis Appl. 7, pp. 268 277 (1973).
- L.A. Belyakov (1): Bifurcation in a system with homoclinic saddle curve. Mat. Zam. 28, pp. 911 - 922 (1980).
- L.A. Belyakov (2): Bifurcation of systems with homoclinic curve of a saddlefocus with saddle quantity zero. Mat. Zam. 36, pp. 838 - 843 (1984).
- M.S. Berger: Nonlinearity and Functional Analysis. Academic Press, New York (1977).

- R.I. Bogdanov (1): Bifurcation of the limit cycle of a family of plane vector fields (russ.). Trudy Sem. I.G. Petrovskogo 2, pp. 23 – 36 (1976); (engl.) Sel. Mat. Sov. 1, pp. 373 – 387 (1981).
- R.I. Bogdanov (2): Versal deformation of a singularity of a vector field on the plane in the case of zero eigenvalues (russ.). Ibid., 37 65; (engl.) Sel. Mat. Sov. 1, pp. 389 421 (1981).
- R.I. Bogdanov (3): Invariants of singular points in the plane (russ.). Uspekhi Mat. Nauk 40, pp. 199 – 200 (1985).
- P. Brunovský & B. Fiedler (1): Numbers of zeros on invariant manifolds in reaction-diffusion equations. Nonl. Analysis TMA 10, pp. 179 193 (1986).
- P. Brunovský & B. Fiedler (2): Connecting orbits in scalar reaction diffusion equations. In: Dynamics Reported 1, U. Kirchgraber & H.-O. Walther (eds.), pp. 57 - 89.
- P. Brunovský & B. Fiedler (3): Connecting orbits in scalar reaction diffusion equations II: the complete solution. To appear in J. Diff. Eq.
- V.V. Bykov: Bifurcations of dynamical systems close to systems with a separatrix contour containing a saddle-focus (russ.). In: Methods of the Qualitative Theory of Differential Equations, E.A. Leontovich-Andronova (ed.), Gor' kov. Gos. Univ., Gorki, pp. 44 - 72 (1980).
- S.-N. Chow & B. Deng: Homoclinic and heteroclinic bifurcation in Banach spaces. Preprint (1984).
- S.-N. Chow & B. Deng & D. Terman (1): The bifurcation of a homoclinic orbit from two heteroclinic orbit a topological approach. Preprint (1986).
- S.-N. Chow & B. Deng & D. Terman (2): The bifurcation of a homoclinic and periodic orbits from two heteroclinic orbits — an analytic approach. Preprint (1987).
- S.-N. Chow & J.K. Hale: Methods of Bifurcation Theory. Grundl. math. Wiss. 251, Springer-Verlag, New York (1982).
- S.-N. Chow & X.-B. Lin: Bifurcation of a homoclinic orbit with a saddle-node equilibrium. Preprint (1988).
- S.-N. Chow & J. Mallet-Paret & J.A. Yorke: A periodic orbit index which is a bifurcation invariant. In: [PALIS], pp. 109 – 131.
- P. Coullet & J.-M. Gambaudo & C. Tresser: Une nouvelle bifurcation de codimension 2: le collage de cycles. C.R. Acad. Sc. Paris 299, pp. 253 – 256 (1984).

- **B.** Deng (1): Šil'nikov problem, exponential expansion, strong λ -Lemma, C^1 -linearization, and homoclinic bifurcation. Preprint (1987).
- B. Deng (2): The bifurcation of a nondegenerate heteroclinic loop with an application to the existence of infinite transition waves in a bistable reaction diffusion system. Preprint (1987).
- **B.** Deng (3): Exponential expansion in Sil'nikov's saddle-focus. Preprint (1988).
- **B. Deng (4)**: Homoclinic bifurcations with nonhyperbolic equilibria. Preprint (1988).
- P. Deufihard & B. Fiedler & P. Kunkel: Efficient numerical pathfollowing beyond critical points. SIAM J. Numer. Analysis 24, pp. 912 - 927 (1987).
- E.J. Doedel & J.P. Kernevez: Software for continuation problems in ordinary differential equations with applications. CALTECH (1985).
- J. Evans & N. Fenichel & J.A. Feroe: Double impulse solutions in nerve axon equations. SIAM J. Appl. Math. 42, pp. 219 234 (1982).
- J.A. Feroe: Existence and stability of multiple impulse solutions of a nerve axon equation. SIAM J. Appl. Math. 42, pp. 235 246 (1982).
- B. Fiedler (1): An index for global Hopf bifurcation in parabolic systems. J. reine angew. Math. 359, pp. 1 36 (1985).
- B. Fiedler (2): Global Hopf bifurcation of two-parameter flows. Arch. Rational Mech. Analysis 94, pp. 59 - 81 (1986).
- B. Fiedler & P. Kunkel (1): A quick multiparameter test for periodic solutions.
 In: Bifurcation: Analysis, Algorithms, Applications, T. Küpper & R. Seydel & H. Troger (eds.), ISNM 79, Birkhäuser, Basel, pp. 61 – 70 (1970).
- B. Fiedler & P. Kunkel (2): Multistability, scaling, and oscillations. In: Complex Chemical Reaction Systems, J. Warnatz & W. Jäger (eds.), Springer Series in Chem. Physics 47, Berlin, pp. 169 180 (1987).
- B. Fiedler & J. Mallet-Paret: Connections between Morse sets for delaydifferential equations. To appear in J. reine angew. Math.
- G. Fischer: Zentrumsmannigfaltigkeiten bei elliptischen Differentialgleichungen. Math. Nachr. 115, pp. 137 – 157 (1984).
- R. FitzHugh: Mathematical models of excitation and propagation in nerves. In: Biological Engineering, H.P. Schwan (ed.), McGraw-Hill, New York, pp. 1 – 85 (1969).

- J.-M. Gambaudo & P. Glendinning & C. Tresser: Collage de cycles et suites de Farey. C. R. Acad. Sc. Paris 299, pp. 711 714 (1984).
- P. Glendinning (1): Bifurcation near homoclinic orbits with symmetry. Phys. Lett. 103 A, pp. 163 166 (1984).
- P. Glendinning (2): Travelling wave solutions near isolated double-pulse solitary waves of nerve axon equations. Phys. Lett. 121 A, pp. 411 – 413 (1987).
- P. Glendinning & C. Sparrow: T-points: a codimension two heteroclinic bifurcation. J. Stat. Physics 43, pp. 479 - 488 (1986).
- J. Guckenheimer & P. Holmes: Structurally stable heteroclinic cycles. Math. Proc. Camb. Phil. Soc. 103, pp. 189 – 192 (1988).
- J.K. Hale & X.-B. Lin: Heteroclinic orbits for retarded functional differential equations. J. Diff. Eq. 65, 175 202 (1986).
- S.P. Hastings: Single and multiple pulse waves for the FitzHugh-Nagumo equations. SIAM J. Appl. Math 42, pp. 247 - 260 (1982).
- M.W. Hirsch & C.C. Pugh & M. Shub: Invariant Manifolds. Lect. Notes Math. 583, Springer-Verlag, Berlin (1977).
- H. Hofer & J. Toland: Homoclinic, heteroclinic, and periodic orbits for a class of indefinite Hamiltonian systems. Math. Ann. 268, pp. 387 403 (1984).
- P. Holmes: A strange family of three-dimensional vector fields near a degenerate singularity. J. Diff. Eq. 37, pp. 382 403 (1980).
- P. de Hoog & R. Weiss: The numerical solution of boundary value problems with an essential singularity. SIAM J. Numer. Analysis 16, 637 - 669 (1979).
- W. Hurewicz & H. Wallman: Dimension Theory. Princeton Univ. Press (1948).
- H. Kielhöfer & R. Lauterbach: On the principle of reduced stability. J. Funct. Analysis 53, pp. 99 - 111 (1983).
- K. Kirchgässner (1): Wave-solutions of reversible systems and applications. J. Diff. Eq. 45, pp. 113 127 (1982).
- K. Kirchgässner (2): Homoclinic bifurcation of perturbed reversible systems. In: Equadiff 82, H.W. Knobloch & K. Schmitt (eds.), Lect. Notes Math. 1017, Springer-Verlag, Heidelberg, pp. 328 - 363 (1983).
- K. Kirchgässner (3): Nonlinearly resonant surface waves and homoclinic bifurcation. Preprint (1988).

- H. Kokubu (1): On a codimension 2 bifurcation of homoclinic orbits. Proc. Japan Acad. 63 A, pp. 298 - 301 (1987).
- H. Kokubu (2): Homoclinic and heteroclinic bifurcations of vector fields. Japan J. Appl. Math. 5, pp. 455 - 501 (1988).
- M. Kubiček & M. Marek: Computational Methods in Bifurcation Theory and Dissipative Structures. Springer-Verlag, New York (1983).
- I. Kupka: Contribution à la théorie des champs génériques. Contrib. Diff. Eq. 2, pp. 457 484 (1963) and 3, pp. 411 420 (1964).
- Y. Kuramoto & S. Koga: Anomalous period-doubling bifurcations leading to chemical turbulence. Physics Letters 92 A, pp. 1 4 (1982).
- M. Lentini & H.B. Keller: Boundary value problems on semi-infinite intervals and their numerical solution. SIAM J. Numer. Analysis 17, pp. 577 - 604 (1980).
- E. Leontovich: On the generation of limit cycles from separatrices (russ.). Dokl. Akad. Nauk 78, pp. 641 – 644 (1951).
- X.-B. Lin: Exponential dichotomies and homoclinic orbits in functional differential equations. J. Diff. Eq. 63, pp. 227 – 254 (1986).
- V.I. Lukyanov: Bifurcations of dynamical systems with a saddle-point separatrix loop. Diff. Eq. 18, pp. 1049 1059 (1982).
- D.V. Lyubimov & M.A. Zaks: Two mechanisms of the transition to chaos in finite-dimensional models of convection. Physica 9D, pp. 52 64 (1983).
- J. Mallet-Paret & J.A. Yorke (1): Two types of Hopf bifurcation points: sources and sinks of families of periodic orbits. In: Nonlinear Dynamics, R.H.G. Helleman (ed.), Proc. NY Acad. Sc. 357, pp. 300 - 304 (1980).
- J. Mallet-Paret & J.A. Yorke (2): Snakes: oriented families of periodic orbits, their sources, sinks, and continuation. J. Diff. Eq. 43, pp. 419 – 450 (1982).
- A. Mielke (1): A reduction principle for nonautonomous systems in infinitedimensional spaces. J. Diff. Eq. 65, pp. 68 - 88 (1986).
- A. Mielke (2): Steady flows of inviscid fluids under localized perturbations. Ibid., pp. 89 116.
- J. Moser: Stable and Random Motions in Dynamical Systems. Princeton University Press (1973).

- V.P. Nozdracheva: Bifurcation of a noncoarse separatrix loop. Differential Eq. 18, pp. 1098 1104 (1982).
- I.M. Ovsyannikov & L.P. Shil'nikov: On systems with a saddle-focus homoclinic curve. Math. USSR Sbornik 58, pp.. 557 - 574 (1987).
- J. Palis jr. (ed.): Geometric Dynamics. Lect. Notes Math 1007, Springer-Verlag, New York (1983).
- K.J. Palmer: Exponential dichotomies and transversal homoclinic points. J. Diff. Eq. 55, pp. 225 - 256 (1984).
- P.H. Rabinowitz: Some global results for nonlinear eigenvalue problems. J. Funct. Analysis 7, pp. 487 513 (1971).
- J.W. Reyn: Generation of limit cycles from separatrix polygons in the phase plane. In: Geometrical Approaches to Differential Equations, R. Martini (ed.), Lect. Notes Math. 810, Springer-Verlag, Berlin, pp. 264 – 289 (1980).
- J. Rinzel & D. Terman: Propagation phenomena in a bistable reaction-diffusion system. SIAM J. Appl. Math 42, pp. 1111 - 1137 (1982).
- C. Robinson: Differentiability of the stable foliation for the model Lorenz equations. Preprint (1988).
- **J.A. Rodriguez:** Bifurcations to homoclinic connections of the focus-saddle type. Arch. Rational Mech. Analysis 93, pp. 81 – 90 (1986).
- J.A. Sanders & R. Cushman: Limit cycles in the Josephson equation. SIAM J. Math. Analysis 17, pp. 495 511 (1986).
- S. Schecter (1): The saddle-node separatrix-loop bifurcation. SIAM J. Math. Analysis 18, pp. 1142 - 1156 (1987).
- S. Schecter (2): Melnikov's method at a saddle-node and the dynamics of the forced Josephson junction. SIAM J. Math. Analysis 18, pp. 1699 1715 (1987).
- G.R. Sell (1): Smooth linearization near a fixed point. Amer. J. Math. 107, pp. 1035 1091 (1985).
- G.R. Sell (2): Obstacles to linearization. Differential Eq. 20, pp. 341 345 (1984).
- R. Seydel: From Equilibrium to Chaos. Elsevier, New York (1988).
- L.P. Shil'nikov (1): Some cases of generation of periodic motions in an n-dimensional space. Soviet Math. Dokl. 3, pp. 394 - 397 (1962).

- L.P. Shil'nikov (2): A case of the existence of a countable number of periodic motions. Soviet Math. Dokl. 6, pp. 163 166 (1965).
- L.P. Shil'nikov (3): On the generation of a periodic motion from a trajectory which leaves and re-enters a saddle-saddle state of equilibrium. Soviet Math. Dokl. 7, pp. 1155 - 1158 (1966).
- L.P. Shil'nikov (4): The existence of a denumerable set of periodic motions in four-dimensional space in an extended neighborhood of a saddle-focus. Soviet Math. Dokl. 8, pp. 54 57 (1967).
- L.P. Shil'nikov (5): On the generation of a periodic motion from trajectories doubly asymptotic to an equilibrium state of saddle type. Math. USSR Sbornik 6, pp. 427 - 437 (1968).
- L.P. Shil'nikov (6): On a new type of bifurcation of multidimensional dynamical systems. Soviet Math. Dokl. 10, pp. 1368 1371 (1969).
- L.P. Shil'nikov (7): A contribution to the problem of the structure of an extended neighborhood of a rough equilibrium state of saddle-focus type. Math. USSR Sbornik 10, pp. 91 – 102 (1970).
- M. Shub: Global Stability of Dynamical Systems. Springer-Verlag, New York (1987).
- S. Smale: Stable manifolds for differential equations and diffeomorphisms. Ann. Sc. Norm. Sup. Pisa 17, pp. 97 116 (1963).
- C. Sparrow: The Lorenz Equations: Bifurcations, Chaos, and Strange Attractors. Appl. Math. Sc. 41, Springer-Verlag, New York (1982).
- S. Sternberg (1): Local contractions and a theorem of Poincaré. Amer. J. Math. 79, pp. 809 824 (1957).
- S. Sternberg (2): On the structure of local homeomorphisms of Euclidean nspace. Amer. J. Math. 80, pp. 623 - 631 (1958).
- F. Takens: Singularities of vector fields. Publ. IHES 43, pp. 47 100 (1974).
- C. Tresser: About some theorems by L.P. Shil'nikov. Ann. Inst. H. Poincaré 40, pp. 441 461 (1984).
- A. Vanderbauwhede: Center manifolds, normal forms and elementary bifurcations. To appear in Dynamics Reported, U. Kirchgraber & H.-O. Walther (eds.).
- H.-O. Walther (1): Homoclinic solution and chaos in $\dot{x}(t) = f(x(t-1))$. Nonl. Analysis TMA 5, pp. 775 – 788 (1981).

- H.-O. Walther (2): Bifurcation from a heteroclinic solution in differential delay equations. Trans. AMS 290, pp. 213 233 (1985).
- H.-O. Walther (3): Inclination lemmas with dominated convergence. J. Appl. Math. Phys. (ZAMP) 32, pp. 327 - 337 (1987).
- H.-O. Walther (4): Bifurcation from a saddle connection in functional differential equations: an approach with inclination lemmas. Preprint (1986).
- H.-O. Walther (5): Homoclinic and periodic solutions of scalar differential delay equations. Preprint (1987).
- E. Yanagida: Branching of double pulse solutions from single pulse solutions in nerve axon equations. J. Diff. Eq. 66, pp. 243 262 (1987).